arXiv:0802.1289v1 [hep-ph] 9 Feb 2008

Preprint typeset in JHEP style - HYPER VERSION

$\pi\eta$ scattering and the resummation of vacuum fluctuation in three-flavour χPT

Marian Kolesár

Institute of Particle and Nuclear Physics, Charles University, V Holešovičkách 2, 180 00
Prague 8, Czech Republic
E-mail: `marian.kolesar@mff.cuni.cz`

Jiří Novotný

Institute of Particle and Nuclear Physics, Charles University, V Holešovičkách 2, 180 00
Prague 8, Czech Republic
E-mail: `jiri.novotny@mff.cuni.cz`

Abstract: We discuss various aspects of resummed chiral perturbation theory, which was developed recently in order to consistently include the possibility of large vacuum fluctuations of the $\bar{s}s$-pairs and the scenario with smaller value of the $\bar{q}q$ condensate for $N_f = 3$. The subtleties of this approach are illustrated using a concrete example of observables connected with $\pi\eta$ scattering. This process seems to be a suitable theoretical laboratory for this purpose due to its sensitivity to the values of the $O(p^4)$ LEC's, namely to the values of the fluctuation parameters L_4 and L_6. We discuss several issues in detail, namely the choice of "good" observables and properties of their bare expansions, the "safe" reparametrization in terms of physical observables, the implementation of exact perturbative unitarity and exact renormalization scale independence, the role of higher order remainders and their estimates. We make a detailed comparison with standard chiral perturbation theory and use generalized χPT as well as resonance chiral theory to estimate the higher order remainders.

Keywords: chiral perturbation theory, resummation of vacuum fluctuations, $\pi\eta$ scattering, resonance chiral theory.

Contents

1. Introduction **2**

2. Resummation of the vacuum fluctuations - motivation and basic notation **6**

3. Construction of the bare expansion for the scattering $P_1P_2 \to P_3P_4$ **8**
- 3.1 Chiral expansion of the “good” observable 8
- 3.2 Dispersive representation for $G(s,t;u)$ 9
- 3.3 Matching the strict chiral expansion to the dispersive representation 12

4. General properties of $\pi\eta$ scattering amplitude **14**
- 4.1 Basic notation 14
- 4.2 Dispersive representation 15
- 4.3 Bare expansion for $G(s,t;u)$ 16

5. Reparametrization of the bare expansion **18**
- 5.1 $\pi\eta$ scattering within the standard chiral perturbation theory to $O(p^4)$ 18
- 5.2 Resummation of the vacuum fluctuation 20
- 5.3 $\pi\eta$ scattering within the generalized chiral perturbation theory to $O(p^4)$ - the bare expansion of $G(s,t;u)$ 21
- 5.4 Observables of the $\pi\eta$ scattering within $G\chi PT$ to $O(p^4)$ - the reparametrization 23

6. Numerical results **25**
- 6.1 The standard chiral perturbation theory 26
- 6.2 Resummation of vacuum fluctuations - basic properties 28
- 6.3 Numerical comparison of the resummed and standard reparametrization 30
- 6.4 The role of the remainders 34
- 6.5 Resonances estimate of the direct remainders 34
- 6.6 Generalized χPT 37

7. Summary and conclusions **39**

A. Standard chiral expansion of the parameters X, Z and r **41**

B. Chiral expansion of the η decay constant **41**

C. Dispersion representation of the $\pi\eta$ amplitude **43**
- C.1 $S-$channel discontinuities at $O(p^4)$ 44
- C.2 $T-$channel discontinuities at $O(p^4)$ 44

D. The scalar bubble **45**

E. $O(p^4)$ constants L_4-L_8 in terms of masses and decay constants **46**

F. Lagrangian of $G\chi PT$ to $O(p^4)$ **46**

G. Coefficients of the dispersive part of $G\chi PT$ amplitude **49**

H. Parameters $\alpha - \omega$ within the generalized χPT **49**

I. Generalized χPT contributions to the bare expansion remainders for the masses 51

J. Resonance amplitude and remainders estimates 53

1. Introduction

As it is well known, at the energy scales $E \ll \Lambda_H \sim 1\text{GeV}$ the physics of QCD is nonperturbative and governed by chiral symmetry (χS) $SU(N_f)_L{\times}SU(N_f)_R$. This global symmetry is present on the classical level within the QCD with N_f massless quarks (in the chiral limit of QCD) and on the quantum level there exist strong theoretical (for $N_f \geq 3$) and phenomenological arguments for spontaneous symmetry breakdown (SSB) of χS according to the pattern $SU(N_f)_L{\times}SU(N_f)_R \to SU(N_f)_V$. Due to the confinement, quark and gluon fields do not represent appropriate low energy degrees of freedom within the above mentioned energy range; the relevant degrees of freedom correspond to the lightest colourless hadrons in the QCD spectrum. As far as the Green functions of quark currents are concerned, it is possible to obtain a general solution of the chiral Ward identities in terms of the low energy expansion. This expansion can be organized most efficiently using the methods of effective field theory corresponding to the low-energy limit of QCD with N_f light quark flavours which is known as chiral perturbation theory (χPT) [1, 2, 3]. χPT describes the low energy QCD dynamics in terms of the lightest $(N_f^2{-}1)$-plet of the pseudoscalar mesons identified with the Goldstone bosons (GB) of the spontaneously broken chiral symmetry which appear in the particle spectrum of the theory as a consequence of the Goldstone theorem. In the chiral limit these pseudoscalars are massless and dominate the low energy dynamics of QCD. They interact weekly at low energies $E \ll \Lambda_H$, where $\Lambda_H \sim 1\text{GeV}$ is the hadronic scale corresponding to the masses of the lightest nongoldstone hadrons. This feature of the GB dynamics enables systematic perturbative treatment with the expansion parameter (E/Λ_H). Within the real QCD the quark mass term $\mathcal{L}^{QCD}_{f,mass}$ breaks χS explicitly and the Goldstone bosons become pseudogoldstone bosons (PGB) with nonzero masses. Though $m_f \neq 0$, for $m_f \ll \Lambda_H$ the mass term $\mathcal{L}^{QCD}_{f,mass}$ can be treated as a perturbation. As a consequence, PGB correspond to the lightest hadrons in the QCD spectrum[1] (identified with $\pi^0, \pi^\pm$ for $N_f = 2$ and $\pi^0, \pi^\pm, K^0, \overline{K^0}, K^\pm, \eta$ for $N_f = 3$) and the interaction of PGB at the energy scale $E \ll \Lambda_H$ continues to be weak. Because $M_P < \Lambda_H$, the QCD dynamics at $E \ll \Lambda_H$ is still dominated by these particles and the effective theory provides us with a simultaneous expansion in powers of (E/Λ_H) and (m_f/Λ_H). The Lagrangian of χPT can be constructed on the basis of symmetry arguments only; the unknown information about the nonperturbative properties of QCD are hidden in the parameters known as low energy constants (LEC)[2, 3]. These are related to the (generally nonlocal) order parameters of the SSB of χS, the most prominent of them are the Goldstone boson decay constant F_0 and the chiral condensate[2] $B_0 = \Sigma/F_0^2$ where $\Sigma = -\langle\overline{u}u\rangle_0$.

To be more precise, N_f-flavour χPT is in fact an expansion in m_i, around the $SU(N_f)_L \times SU(N_f)_R$ chiral limit $m_i = 0$, $i \leq N_f$, while keeping all the other quark masses for $i > N_f$ at their physical values. Because $m_{u,d}$ are much smaller not only in comparison with the hadronic scale

---

[1] The PGB masses M_P can be expanded in the powers (and logarithms) of the quark masses starting from the linear term and therefore vanish in the chiral limit.

[2] The parameter F_0 is however more fundamental in the sense that $F_0 \neq 0$ is both necessary and sufficient condition for SSB, while $\langle\overline{q_f}q_f\rangle_0 \neq 0$ corresponds to the sufficient condition only. (The lower index zero means here the chiral limit.)

Λ_H, but also in comparison with the intrinsic QCD scale Λ_{QCD}, the two-flavour χPT is expected to produce well-behaved expansion corresponding to small corrections to the $SU(2)_L \times SU(2)_R$ chiral limit.

The strange quark mass on the other hand, though still small enough with respect to Λ_H to be treated as an expansion parameter within the three-flavour χPT (relating real QCD with its $SU(3)_L \times SU(3)_R$ chiral limit), is of comparable size with respect to Λ_{QCD}. This fact, besides the expected worse convergence of the three-flavour χPT, might also have interesting consequences for the possible difference between the $N_f = 2$ and $N_f = 3$ chiral dynamics. As discussed intensively in a series of papers [4, 5, 6, 7, 8, 9], $m_s \lesssim \Lambda_{QCD}$ suggest, that the loop effects of the vacuum $\overline{s}s$ pairs are not suppressed as strongly as it is for the heavy quarks and might enhance the magnitude of the $N_f = 2$ chiral order parameters relatively to their $N_f = 3$ chiral limits. This applies mainly to $F_0(N_f)$ and $\Sigma(N_f) = F_0^2(N_f)B_0(N_f)$, which should satisfy paramagnetic inequalities [4]

$$\begin{aligned} \Sigma(2) &> \Sigma(3) = \lim_{m_s \to 0} \Sigma(2) \\ F_0(2) &> F_0(3) = \lim_{m_s \to 0} F_0(2). \end{aligned} \tag{1.1}$$

The leading order difference between the two-flavour and three-flavour values is proportional to m_s, with coefficients measuring the violation of the OZI rule in the 0^{++} channel, *e.g.*

$$\Sigma(2) = \Sigma(3) + m_s \overline{Z}_1^s + \ldots \tag{1.2}$$

(see [4] for details) where

$$\overline{Z}_1^s = \lim_{m_s \to 0} \int \mathrm{d}^4 x \langle \overline{u}u(x) \overline{s}s(0) \rangle_c \tag{1.3}$$

and analogously for F_0. The fluctuation parameter $\overline{Z}_1^s$ is related to the LEC $L_6^r(\mu)$ (and $L_4^r(\mu)$ for F_0) of the three-flavour χPT . As discussed in [4, 5, 6, 7, 8, 9], these parameters might be larger than their estimate based on the large N_c expansion, provided $N_f = 3$ is close to the critical number of light quark flavours N_f^{crit}, for which the chiral symmetry is restored. Available estimates vary widely, some indicate a larger number $N_f^{crit} \sim 10 - 12$ for $N_c = 3$ [10, 11, 12], while other approaches [13, 14] and lattice calculations [15, 16, 17, 18, 19] discuss a possibly much lower value $N_f^{crit} \leq 6$. Provided the scenario of large vacuum fluctuations takes place, the second term in (1.2) (called the induced condensate in [5, 8, 20]) can be numerically comparable with the first term and the three-flavour condensate $\Sigma(3)$ could be substantially smaller than the two-flavour one, the value of which is experimentally accessible in the recent experiments. Analogous reasonings apply to the relationship of $F_0(2)$ and $F_0(3)$.

These effects could possibly have strong consequences for the organization of the chiral expansion in the $N_f = 3$ case [4, 6, 7, 8]. Let us remind that the general form of the Lagrangian of χPT is

$$\mathcal{L} = \sum_{m,n} \mathcal{L}^{(m,n)} \tag{1.4}$$

where

$$\mathcal{L}^{(m,n)} = \sum_k C_k^{(m,n)} O_k^{(m,n)}. \tag{1.5}$$

with LEC's $C_k^{(m,n)}$ and independent set of the operators $O_k^{(m,n)} = O(\partial^m m_f^n)$.

In order to be able to treat the double expansion consistently, it is necessary to assign a single integer parameter called *chiral order* to each term $\mathcal{L}^{(m,n)} = O(\partial^m m_f^n)$ of the effective Lagrangian. The terms $\mathcal{L}_k$ with chiral order k are then called $O(p^k)$ terms. Obviously, $\partial = O(p)$. The matter of discussion might be, however, the question concerning the chiral power of m_f. This question is intimately connected to the scenario according which the SSB of χS is realized.

The standard scenario [2, 3] corresponds to the assumption, that the SSB order parameters $\Sigma(N_f)$ and $F_0(N_f)$ are large in the sense, that the ratios

$$X(N_f) = \frac{2\widehat{m}\Sigma(N_f)}{F_\pi^2 M_\pi^2} \tag{1.6}$$

(where and $\widehat{m} = (m_u + m_d)/2$) and

$$Z(N_f) = \frac{F_0^2(N_f)}{F_\pi^2} \tag{1.7}$$

are close to one. Because $M_\pi^2 = O(p^2)$, it is then natural to take $m_f = O(p^2)$, i.e. $k = m + 2n$. This results in the *standard* χPT ($S\chi PT$ in what follows). This scenario seems to be experimentally confirmed [21] for $N_f = 2$; the recent analysis of the data yields [22]

$$X(2) = 0.81 \pm 0.07, \quad Z(2) = 0.89 \pm 0.03. \tag{1.8}$$

The $O(p^2)$ Lagrangian [2, 3]

$$\mathcal{L}_2 = \frac{F_0^2}{4}\left(\langle \partial_\mu U^+ \partial^\mu U\rangle + 2B_0\langle U^+\mathcal{M} + \mathcal{M}^+ U\rangle\right) \tag{1.9}$$

gives $\Sigma(2)_{LO} = \Sigma(3) = B_0 F_0^2$ at the leading order, thus postponing the difference $\Sigma(2)-\ \Sigma(3)$ to higher orders. The same is true for the parameters $F_0(N_f)$. Let us also note that the quark mass ratio $r = m_s/\widehat{m}$ is not a free parameter here[3], at the leading order one has

$$r = 2\frac{M_K^2}{M_\pi^2} - 1. \tag{1.10}$$

An alternative way of chiral power counting for $N_f = 3$ is the *generalized* χPT ($G\chi PT$) [24, 25, 26, 27, 28, 29], originally designed to treat the scenario with small quark condensate $X(3) \ll 1$ and to take the quark mass ratio r as a free parameter. In the case $X(3) \ll 1$ it is natural to take $m_f = O(p)$ and $B_0 = O(p)$, this means $k = m + n$. In contrast to $S\chi PT$, there are also odd chiral orders and the $O(p^2)$ Lagrangian contains additional terms which are $O(p^4)$ within the standard chiral counting[4](see *e.g.* [24, 25, 28]):

$$\begin{aligned}\mathcal{L}_2 \;=\; & \frac{F_0^2}{4}\big(\langle \partial_\mu U^+ \partial^\mu U\rangle + 2B_0\langle U^+\mathcal{M} + \mathcal{M}^+ U\rangle + A_0\langle (U^+\mathcal{M})^2 + (\mathcal{M}^+ U)^2\rangle \\ & + Z_0^P\langle U^+\mathcal{M} - \mathcal{M}^+ U\rangle^2 + Z_0^S\langle U^+\mathcal{M} + \mathcal{M}^+ U\rangle^2\big)\,. \end{aligned} \tag{1.11}$$

For the condensate $\Sigma = -\langle \overline{u}u\rangle$ we get at the leading order for $N_f = 3$

$$\Sigma_{LO} = B_0 F_0^2 + Z_0^S(2\widehat{m} + m_s) = \Sigma(3) + Z_0^S(2\widehat{m} + m_s) \tag{1.12}$$

and therefore

$$\Sigma_{LO}(2) = \Sigma(3) + Z_0^S m_s. \tag{1.13}$$

This allows the difference $\Sigma(2)-\ \Sigma(3)$ to appear already at the leading order, consistently with the small $\Sigma(3)$ scenario. The next-to-leading order Lagrangian $O(p^3)$

$$\mathcal{L}_3 = \frac{F_0^2}{4}\left(\xi\langle \partial_\mu U^+ \partial^\mu U U^+\mathcal{M} + \mathcal{M}^+ U\rangle + \widetilde{\xi}\langle \partial_\mu U^+ \partial^\mu U\rangle\langle U^+\mathcal{M} + \mathcal{M}^+ U\rangle + \ldots\right) \tag{1.14}$$

---

[3] On the contrary, the value of r is usually taken as an input in standard $O(p^6)$ fits, see *e.g.* [23] and references therein.

[4] Effectively the generalized chiral power-counting means partial resummation of these terms.

(where the ellipses stand for the additional terms which are of the order $O(p^6)$ in $S\chi PT$) gives rise to the $N_f = 3$ relation

$$F^2_{\pi,NLO} = F^2_0(3)(1 + 2\widetilde{\xi}(m_s + 2\widehat{m}) + 2\widehat{m}\xi), \tag{1.15}$$

which implies that the difference $F^2_0(2) - F^2_0(3)$ is treated as an effect of the next-to-leading order

$$F^2_0(2) = F^2_0(3)(1 + 2\widetilde{\xi}m_s). \tag{1.16}$$

Therefore, neither $S\chi PT$ nor $G\chi PT$ can accumulate the case of large fluctuation parameter $\widetilde{\xi}$ and the ratio $Z(3) \ll 1$ at the leading order.

Quite recently, a consistent method of handling the case $X(3)$, $Z(3) \ll 1$ was proposed [4, 6, 7, 8, 9]. Instead of changing the chiral power counting, it is based on a more careful manipulations with the chiral expansion. As it was discussed in the above references, the case $X(3)$, $Z(3) \ll 1$ could significantly influence the properties of the chiral expansion inducing instabilities of the perturbative series corresponding to the observables, which cannot be linearly related to the QCD correlators (such as the *ratios* like PGB masses, scattering amplitudes etc.). For such quantities, one should not perform a perturbative chiral expansion of the denominators but rather keep the ratios in a nonperturbative "resummed" form. The possibly large vacuum $\overline{s}s$ pair fluctuations are then parameterized in terms of $X(3)$, $Z(3)$ and r and treated as free parameters. We return to the detailed formulation of this recipe in the next section.

The aim of this paper is to illustrate the "resummed" form of the chiral expansion with special attention to its formal properties and to the details and subtleties of the general procedure. Motivated by our preliminary results on $\pi\eta$ scattering within the $G\chi PT$ [34], we have chosen the observables connected with this process as a concrete example which seems to be sensitive to the deviations from the standard assumption $X(3)$, $Z(3) \sim 1$ (note that some recent phenomenological studies suggest a possibility of $X(3) \sim 0.5$, cf. [30, 31, 32] and [6, 33]). Also, from the phenomenological point of view, the off-shell $\pi\eta\pi\eta^*$ vertex is a necessary building block for the non-resonant part of the amplitude for the rare decay $\eta \to \pi^0\pi^0\gamma\gamma$. Preliminary estimates within $G\chi PT$ [35, 36] suggest, that the effect of deviation of this off-shell vertex from the standard case might be at least in principle observed. The details will be presented elsewhere [37].

The amplitude of $\pi\eta$ scattering was already calculated within $S\chi PT$ to $O(p^4)$ (and within the extended $S\chi PT$ with explicit resonance fields) in the paper [38], where the authors presented prediction for the scattering lengths and phase shifts of the S, P and D partial waves. We quote here their $O(p^4)$ results for the S- and P-wave scattering lengths (in the units of the pion Compton wavelength): $a_0^{S\chi PT} = 7.2 \times 10^{-3}$ and $a_1^{S\chi PT} = -5.2 \times 10^{-4}$.

The paper is organized as follows. In Section 2 we recapitulate the motivation for the resummed version of χPT and the construction of the bare expansion of "good" observables. We make a detailed general discussion, connected with the four-meson amplitude, of the strict chiral expansion, the dispersive representation and matching of both approaches with stress to the reconciling of exact perturbative unitarity and exact renormalization scale independence in Section 3. Section 4 is devoted to the general properties of the $\pi\eta$ scattering amplitude. We discuss the kinematics, the definition of suitable "good" observables, the dispersion representation of the amplitude and the construction of the bare expansion. Various possibilities of its reparametrization are described in a detailed way in Section 5. The numerical illustration of the particular variants is made in Section 6, where we also numerically illustrate the subtleties of the construction of the bare expansion. We recapitulate the results of the standard variant of χPT and compare them with the resummed approach. We concentrate on the dependence on the LEC's as well as on the sensitivity to the higher order reminders and make an attempt to estimate their values using a matching with $G\chi PT$ and a simple version of resonance chiral theory. Section 7 contains the summary and conclusions. Some technical details are postponed to the appendices.

2. Resummation of the vacuum fluctuations - motivation and basic notation

As we have mentioned in the Introduction, the potentially large vacuum fluctuations of the $\bar{s}s$ pairs might result in the instabilities of the chiral expansion, which originate in the possibility that for some observables the next-to-leading order correction could be numerically comparable with the leading order one. As discussed in [8, 9], this could generally cause problems with the convergence of the formal chiral expansion. Nevertheless, at least for some carefully defined "good" observables, it is natural to presume some sort of satisfactory convergence properties. Such "good" observables are assumed to be those which can be obtained directly from the low-energy correlation functions in the domain of their analyticity far away from singularities and which are related to the corresponding correlator *linearly* [8, 9]. Typical examples are the squares of the PGB decay constants F_P^2 , the products $F_P^2 M_P^2$ where M_P are the PGB masses and also the subthreshold parameters which can be derived from the products $A\prod_{i=1}^{4} F_{P_i}$ where A is the PGB scattering amplitude $1+2 \to 3+4$. Let us write the expansion of such a "good" observable G in the form of a (carefully defined) *bare expansion* [8] as[5]

$$G = G^{(2)} + G^{(4)} + G\delta_G, \tag{2.1}$$

where $G^{(2)} = g^{(2)}(F_0, B_0, m_q)$ and $G^{(4)} = g^{(4)}(F_0, B_0, m_q, L_i, M_P^2)$ correspond to the sum of the leading and next-to-leading order terms respectively and the renormalization scale independent quantity δ_G accommodates the higher order remainders.

As a terminological note, in what follows we use the term *strict* chiral expansion for an unmodified expansion in terms of the LEC's strictly respecting the chiral orders. The *bare* expansion, though still expressed in terms of LEC's, accumulates some modifications dictated by physical requirements. It is the bare expansion which is assumed to be globally convergent.

For a "good" observable it is then assumed

$$|\delta_G| \ll 1 \tag{2.2}$$

as a natural assumption. This property of the bare chiral expansion (2.1) is called global convergence in [8, 9]. Note however, that the validity of the inequality (2.2) might depend on the definition of the reminder δ_G which is not fixed unambiguously and might differ according to the calculation scheme in use. We will comment on this point later on.

The above mentioned possible instability in (2.1) appears when $G^{(2)} \sim G^{(4)}$, *i.e.* $X_G \lessapprox 1$, where

$$X_G = \frac{G^{(2)}}{G}. \tag{2.3}$$

Such an instability manifests itself in the expansion of the observables depending on G nonlinearly [8]. For instance, for a ratio of two "good" observables G and $G^{'}$ formally expanded in the form (2.1)

$$\frac{G}{G'} = \left(\frac{G^{(2)}}{G'^{(2)}}\right) + \left(\frac{G^{(2)}}{G'^{(2)}}\right)\left(\frac{G^{(4)}}{G^{(2)}} - \frac{G'^{(4)}}{G'^{(2)}}\right) + \frac{G}{G'}\delta_{G/G'}, \tag{2.4}$$

we get for the remainder $\delta_{G/G'}$

$$\delta_{G/G'} = \frac{(1-X_{G'})(X_G - X_{G'})}{X_G'^2} + \frac{\delta_G}{X_{G'}} - \frac{X_G \delta_{G'}}{X_G'^2}. \tag{2.5}$$

For $X_{G'} \lessapprox 1$ this might be numerically large even if both $|\delta_G|$, $|\delta_{G'}|$ were reasonably small. In this sense, a ratio of two globally convergent observables need not to be necessarily globally convergent too. It should be therefore much safer not to expand such “dangerous” observables and rather write the ratio in the “resummed” form

---

[5] Here we tacitly assume the standard chiral power counting. Analogous expansion could be written also for the generalized case.

$$\frac{G}{G'} = \frac{G^{(2)} + G^{(4)}}{G'^{(2)} + G'^{(4)}} + \frac{G}{G'}\widetilde{\delta}_{G/G'}. \tag{2.6}$$

The relation (2.6) is an exact algebraic identity provided we keep explicitly the remainder

$$\widetilde{\delta}_{G/G'} = \frac{\delta_G - \delta_{G'}}{1 - \delta_{G'}}. \tag{2.7}$$

In this case $\widetilde{\delta}_{G/G'}$ remains for $|\delta_G|, |\delta_{G'}| \ll 1$ under numerical control.

Of course, only the fact that the bare expansion of some observable is not globally convergent does not necessarily correspond to the collapse of the convergence, because the next-to-next-to-leading order $G^{(6)}$ can saturate the series in such a way that the next-to-next-to-leading remainder

$$G\delta_G^{NNLO} = G - G^{(2)} - G^{(4)} - G^{(6)} \tag{2.8}$$

is reasonably small. Namely this is the usual assumption behind the $O(p^6)$ calculations. Violation of the global convergence property here means merely that the $O(p^6)$ contribution have unnatural size, *i.e.* $G^{(6)} \lesssim G^{(2)} + G^{(4)}$. This could, however, destabilize the $O(p^6)$ chiral expansion of ratios in the way similar to that discussed above.

Provided we allow the expansion of the “good” observables only, we are also pressed to modify the next step leading from the bare expansion to the usual output of the χPT, consisting of a reparametrization of the expansion by expressing some of the LEC's in terms of the physical observables such as masses and PGB decay constants. This step converts the series into an expansion in powers and logs of the (squared) PGB masses instead of quark masses. To achieve this, it is either necessary to invert a bare chiral expansion of some observable (in the case of the $O(p^2)$ LEC's) or to use an observable which might be generally a “dangerous” one. Let us briefly discuss the first case. Schematically, suppose that some $O(p^2)$ LEC G_0 (*e.g.* F_0^2) just corresponds to the leading term $G^{(2)}$ of the expansion of the observable G. Then we can write an algebraic identity

$$G_0 = G - G^{(4)}(G_0) - G\delta_G, \tag{2.9}$$

where we explicitly point out the dependence of the next-to-leading term on G_0. To convert this expansion and express G_0 by means of the series in G one substitutes G for G_0 on the right hand side. This defines a new remainder δ_{G_0}

$$G_0 = G - G^{(4)}(G) + G_0\delta_{G_0}, \tag{2.10}$$

for which we get

$$\delta_{G_0} = -\frac{1 - X_G}{X_G} + \frac{1}{X_G}\frac{G^{(4)}(G)}{G}. \tag{2.11}$$

This could cause an instability of the converted expansion for G_0 in terms of G for $X_G \lnsim 1$ even if the relative size of the next-to-leading order $G^{(4)}(G)/G$ is reasonably small, irrespective of the condition for global convergence $|\delta_G| \ll 1$.

On the other hand, suppose that some $O(p^4)$ constant G_1 coincides with the next-to-leading term $G^{(4)}$. In this case we have an algebraic identity for G_1

$$G_1 = G - G^{(2)} - G\delta_G \tag{2.12}$$

and the remainder here is perfectly under control, provided G has a globally convergent bare expansion and we *do not* re-express $G^{(2)}$ in terms of physical observables (i.e. provided we treat the $O(p^2)$ LEC's as free parameters).

From the above simple considerations follows that in order to avoid potential problems with the instabilities of the chiral expansion, which might be present in the three-flavor χPT in the case of small $X(3)$ and $Z(3)$ (cf. (1.6, 1.7)), we should [8, 9]

- carefully define the bare expansion
- confine ourselves (as far as the bare chiral expansion is concerned) to the linear space of "good" observables and keep the "dangerous" observables in the nonperturbative "resummed" form
- use rather $\Sigma(3)$, $F_0(3)$ (or $X(3)$ and $Z(3)$) and $r = 2m_s/(m_u + m_d)$ as free parameters[6] instead of expressing them in the form of the series in PGB masses and decay constants
- eliminate the $O(p^4)$ LEC's algebraically, using bare expansions of "good" observables such as F_P^2, $F_P^2 M_P^2$ [7].

In the next section we shall illustrate the possible subtleties of the first step of this general recipe on the concrete example[8] of the PGB scattering amplitude $P_1P_2 \to P_3P_4$

3. Construction of the bare expansion for the scattering $P_1P_2 \to P_3P_4$

3.1 Chiral expansion of the "good" observable

Let us assume a scattering of pseudoscalar mesons $P_1P_2 \to P_3P_4$ with masses M_{P_i}. The amplitude $S(s,t;u)$ is defined as

$$\langle P_3(k_3)P_4(k_4)_{out}|P_1(k_1)P_2(k_2)_{in}\rangle = \mathrm{i}(2\pi)^4\delta^{(4)}(k_3 + k_4 - k_1 - k_2)S(s,t;u), \tag{3.1}$$

where s, t and u are the usual Mandelstam variables. The amplitude is related to the "good" observable[9]

$$G(s,t;u) = \prod_{i=1}^{4} F_{P_i} S(s,t;u), \tag{3.2}$$

(where F_{P_i} are the decay constants) which can be directly obtained from the (cut) four-point function of the axial currents. Let us write for $G(s,t;u)$ the following *strict* chiral expansion in terms of the low energy constants

$$G = G^{(2)} + G^{(4)}_{ct} + G^{(4)}_{tad} + G^{(4)}_{unit} + G\delta_G. \tag{3.3}$$

$G\delta_G$ accommodates the higher order remainders. Using the functional method, G can be obtained from the generating functional

$$\begin{aligned} F_0^4 Z[U,v,p,a,s] \;=\; & F_0^4 \int d^4x \left(\mathcal{L}^{(2)}(U,v,p,a,s) + \mathcal{L}^{(4)}(U,v,p,a,s)\right) \\ & + F_0^4 Z^{(4)}_{loop}[U,v,p,a,s] + \ldots \end{aligned} \tag{3.4}$$

---

[6] Note, that r is related to the "dangerous" observable

$$2\frac{F_K^2 M_K^2}{F_\pi^2 M_\pi^2} - 1 = r + \ldots$$

[7] We will do it for $L_4 - L_8$ but leave $L_1 - L_3$ free, also L_7 is a special case, see in what follows.

[8] We shall tacitly assume the case of three light flavours in what follows.

[9] Strictly speaking, the "good" observables correspond to the subthreshold parameters derived form $G(t,s;u)$ in an unphysical point away from singularities.

by setting $v = s = p = 0$, $s = 2B_0\mathcal{M}$ and expanding in the fields Φ where $U = \exp(i\Phi/F_0)$. Following the notation in [3], we have

$$\begin{aligned} Z^{(4)}_{loop}[U,v,p,a,s] &= Z^{(4)}_{tad}[U,v,p,a,s] + Z^{(4)}_{unit}[U,v,p,a,s] \\ &= \frac{i}{2}\ln\det D_0 + \frac{i}{4}\mathrm{Tr}(D_0^{-1}\delta) - \frac{i}{4}\mathrm{Tr}(D_0^{-1}\delta D_0^{-1}\delta) + \ldots . \end{aligned} \tag{3.5}$$

In the above formulae,

$$D_0^{ab} = \delta^{ab}\Box + \frac{1}{2}B_0 tr(\{\lambda^a,\lambda^b\}\mathcal{M}) \tag{3.6}$$

and $\mathcal{M}$ is the quark mass matrix. Note this representation of $Z^{(4)}_{loop}$ assumes that the masses running in the loops are the $O(p^2)$ masses rather than the physical masses. Or, in more detail, provided we start with the chiral expansion of the squared product of the masses and decay constants

$$F_P^2 M_P^2 = (F_P^2 M_P^2)^{(2)} + (F_P^2 M_P^2)^{(4)} + F_P^2 M_P^2 \delta_{FM_P}, \tag{3.7}$$

the masses in the loops are defined as

$$\overset{o}{M}{}_P^{2} = \frac{(F_P^2 M_P^2)^{(2)}}{F_0^2}. \tag{3.8}$$

Note, however, that this is the first term in a potentially "dangerous" expansion of the ratio

$$M_P^2 = \frac{F_P^2 M_P^2}{F_P^2} = \frac{(F_P^2 M_P^2)^{(2)} + (F_P^2 M_P^2)^{(4)} + F_P^2 M_P^2 \delta_P}{F_0^2 + (F_P^2)^{(4)} + F_P^2 \delta_{F_P}} = \overset{o}{M}{}_P^{2} + \ldots \tag{3.9}$$

From this definition of $Z^{(4)}_{loop}$ we obtain $G^{(4)} = G^{(4)}_{ct} + G^{(4)}_{tad} + G^{(4)}_{unit}$ which is exactly renormalization scale independent even for the external momenta off-shell. This meets the requirement of the renormalization scale independence of the remainder δ_G.

The first two terms of the above strict chiral expansion for $G(s,t;u)$ have a serious drawback in the sense that the singularities in the complex stu planes required by unitarity are not placed at the physical thresholds but rather at points given by the leading order terms $\overset{o}{M}_P$ of the chiral expansion of the PGB masses. Straightforward substitution $\overset{o}{M}_P \rightarrow M_P$ in the propagators of the loops, which apparently means merely a redefinition of the remainder δ_G, could, however, in general spoil its exact renormalization scale independence. It is therefore desirable to use the freedom in the definition of the remainder more carefully in order to reconcile both scale independence of $G^{(4)}$ and unitarity. For this purpose, a useful tool is the matching with a dispersive representation [8] of the amplitude $S(s,t;u)$ based on the reconstruction theorem [25, 29].

3.2 Dispersive representation for $G(s,t;u)$

The above mentioned reconstruction theorem for the PGB scattering amplitude is based on the basic properties of unitarity, analyticity and crossing symmetry and provides us with the most general form of the PGB scattering amplitude up to the order $O(p^6)$ in terms of dispersive integrals with known discontinuities. It was first proved for the case of $\pi\pi$ scattering in [25, 29] and for πK scattering in [39, 32] and since then it has been intensively used in various contexts. Here we use the general form of the theorem, more detailed discussion of which will be presented elsewhere [40].

For the scattering of pseudoscalar mesons $P_1P_2 \rightarrow P_3P_4$, let us denote the $s-$, $t-$ and $u-$channel amplitudes as $S(s,t;u)$, $T(s,t;u)$ and $U(s,t;u)$ and write their partial wave expansion as

$$A(s,t;u) = 32\pi \sum_{l=0}^{\infty}(2l+1)A_l(s)P_l(\cos\theta_A), \tag{3.10}$$

where $A = S$, T, U and

$$\cos\theta_A = \frac{s(t-u) + \Delta_{A_i}\Delta_{A_f}}{\lambda_{A_i}^{1/2}(s)\lambda_{A_f}^{1/2}(s)}. \tag{3.11}$$

Here $A_l(s)$ are the partial waves,

$$\lambda_{A_{i,f}}(s) = (s - (M_{P_j} + M_{P_k})^2)(s - (M_{P_j} - M_{P_k})^2) \tag{3.12}$$

is the triangle function which corresponds to the initial/final state $A_{i,f}$ (consisting of the pseudoscalars P_jP_k) of the process in the channel A and

$$\Delta_{A_{i,f}} = M_{P_j}^2 - M_{P_k}^2. \tag{3.13}$$

According to the theorem, we get the following representation for the amplitude $S(s,t;u)$

$$S(s,t;u) = \mathcal{S}(s,t;u) + \mathcal{S}_{unit}(s,t;u) + O(p^8), \tag{3.14}$$

where $\mathcal{S}(s,t;u)$ is a third order polynomial with the same symmetries as the whole amplitude $S(s,t;u)$. The nontrivial analytical properties are incorporated in the unitarity part $\mathcal{S}_{unit}(s,t;u)$, which can be expressed as

$$\begin{aligned}\mathcal{S}_{unit}(s,t;u) \;=\; & \Phi^S(s) + \Phi^T(t) + \Phi^U(u) \\ & +[s(t-u) + \Delta_{12}\Delta_{34}]\Psi^S(s) \\ & +[t(s-u) + \Delta_{13}\Delta_{24}]\Psi^T(t) \\ & +[u(t-s) + \Delta_{14}\Delta_{23}]\Psi^U(u).\end{aligned} \tag{3.15}$$

In the last expression, $\Delta_{ij} = M_{P_i}^2 - M_{P_j}^2$. The functions $\Phi^A(s)$ and $\Psi^A(s)$ with $A = S,T,U$ are analytic in the cut complex plane with the right hand cut from $\tau_A = \min_{i,j}(M_{P_i} + M_{P_j})^2$ (where P_iP_j are the possible intermediate states in the given channel A) to infinity with discontinuities given by the formulae

$$\mathrm{disc}\,\Phi^A(s) \;=\; 32\pi\theta(s-\tau_A)\mathrm{disc}\,A_0(s) \tag{3.16}$$

$$\mathrm{disc}\,\Psi^A(s) \;=\; 96\pi\theta(s-\tau_A)\mathrm{disc}\frac{A_1(s)}{\lambda_{A_i}^{1/2}(s)\lambda_{A_f}^{1/2}(s)}. \tag{3.17}$$

Here $A_0(s)$, $A_1(s)$ are the corresponding $l = 0, 1$ partial waves.

Consequently, once the right hand sides of (3.16, 3.17) are known, the unitarity part $\mathcal{S}_{unit}(s,t;u)$ of the amplitude can be uniquely reconstructed to $O(p^6)$ up to the polynomial, which encompass subtraction polynomials for the dispersion integrals.

Let us now assume the chiral expansion of the amplitudes in the form

$$A(s,t;u) \;=\; A^{(2)}(s,t;u) + A^{(4)}(s,t;u) + A\delta_A, \tag{3.18}$$

$$A^{(n)}(s,t;u) \;=\; 32\pi\sum_{l=0}^{\infty}(2l+1)A_l^{(n)}(s)P_l(\cos\theta_s(t)). \tag{3.19}$$

Starting from the $O(p^2)$ amplitudes, we can use the two particle partial wave unitarity to get the discontinuity of the partial waves $A_l^{(4)}(s)$ along the right hand cut[10]

$$\mathrm{disc}A_l^{(4)}(s) = \sum_{ij}\frac{2}{z_{ij}}\frac{\lambda_{ij}^{1/2}(s)}{s}A_l^{(2)ij\to A_f}(s)A_l^{(2)ij\to A_i}(s)^* + O(p^6) \tag{3.20}$$

---

[10] It can be shown that more than two particle intermediate states yield contribution of the order $O(p^8)$ and higher.

Here $z_{ij} = 1, 2$ is a symmetry factor taking into account the possibility of identical particles in the intermediate state ij. Inserting this into the dispersive integrals we easily[11] get a minimal form for the $O(p^4)$ unitarity corrections in terms of the functions $\Phi^{(4)A}(s)$ and $\Psi^{(4)A}(s)$ reconstructed from the $O(p^2)$ amplitudes

$$\begin{aligned} \Phi^{(4)A}(s) &= (32\pi)^2 \sum_{ij} \frac{1}{z_{ij}} \overline{\overline{J}}_{ij}(s) A_0^{(2)A_i \to ij}(s) A_0^{(2)ij \to A_f}(s)^* \qquad (3.21) \\ \Psi^{(4)A}(s) &= \frac{(96\pi)^2}{3} \sum_{ij} \frac{1}{z_{ij}} \overline{\overline{J}}_{ij}(s) \frac{A_1^{(2)A_i \to ij}(s) A_1^{(2)ij \to A_f}(s)^*}{\lambda_{A_i}^{1/2}(s) \lambda_{A_f}^{1/2}(s)}. \qquad (3.22) \end{aligned}$$

Provided $\Delta_{A_i} = 0$ or $\Delta_{A_f} = 0$, which will be our case, it can be shown that we only need one subtraction, $\overline{J}_{ij}(s) = J_{ij}^r(s) - J_{ij}^r(0)$ instead of $\overline{\overline{J}}_{ij}(s)$. The explicit form of the function $\overline{J}_{ij}(s)$ is given in the Appendix D.

The above formulae can be used to write a dispersive representation of the "good" observable $G(s, t; u) = \prod_{i=1}^4 F_{P_i} S(s, t; u)$ to the next-to-leading order in the form

$$G(s, t; u) = \mathcal{G}(s, t; u) + \mathcal{G}_{unit}(s, t; u), \qquad (3.23)$$

where $\mathcal{G}(s, t; u)$ is the polynomial part and the unitarity corrections up to $O(p^6)$ are included in

$$\begin{aligned} \mathcal{G}_{unit}(s, t; u) &= \phi^S(s) + \phi^T(t) + \phi^U(u) \\ &\quad + [s(t-u) + \Delta_{12}\Delta_{23}]\psi^S(s) \\ &\quad + [t(s-u) + \Delta_{13}\Delta_{24}]\psi^T(t) \\ &\quad + [u(t-s) + \Delta_{14}\Delta_{23}]\psi^U(u). \qquad (3.24) \end{aligned}$$

Our goal is to write down a representation of $\phi^{(4)A}$ and $\psi^{(4)A}$, which, notice, are distinct quantities from $\Phi^{(4)A}$ and $\Psi^{(4)A}$, analogous to (3.21, 3.22). Note, however, that while the relation of $G(s, t; u)$ and $S(s, t; u)$ is unambiguously fixed to all orders by (3.2), the amplitude can be defined order by order in various ways. For example, for the "good" observable G, the leading order piece $G^{(2)}$ of its strict chiral expansion is fixed by the lowest order Lagrangian $\mathcal{L}^{(2)}$, but the corresponding $O(p^2)$ piece of the amplitude S can be related in various ways. Similarly, the same is true order by order, where the amplitude at the given order can be defined up to higher order corrections.

The most straightforward way is to write a safe expansion for $S(s, t; u)$ in the form

$$S(s, t; u) = \left(\prod_{i=1}^{4} F_{P_i} \right)^{-1} \left(G^{(2)}(s, t; u) + G^{(4)}(s, t; u) + G\delta_G \right) \qquad (3.25)$$

with physical values of F_{P_i}, thus satisfying the relation (3.2) order by order

$$S^{(n)}(s, t; u) = \left(\prod_{i=1}^{4} F_{P_i} \right)^{-1} G^{(n)}(s, t; u). \qquad (3.26)$$

As we'll see, the minimal modification of the form derived from the generating functional is obtained by using an alternative, potentially "dangerous" expansion

---

[11] Note that $A^{(2)}(s, t; u)$ are real polynomials of the first order in s, t and u.

$$\begin{aligned}
S(s,t;u) &= \left(\prod_{i=1}^{4} F_{P_i}\right)^{-1} G(s,t;u) \\
&= \left(\prod_{i=1}^{4} F_0\Big(1+\frac{1}{2}\frac{(F_{P_i}^2)^{(4)}}{F_0^2}+\ldots\Big)\right)^{-1} \Big(G^{(2)}(s,t;u)+G^{(4)}(s,t;u)+\ldots\Big) \\
&= F_0^{-4}G^{(2)}(s,t;u)-\frac{1}{2}F_0^{-6}G^{(2)}(s,t;u)\sum_{i=1}^{4}(F_{P_i}^2)^{(4)}+F_0^{-4}G^{(4)}(s,t;u)+\ldots, \qquad (3.27)
\end{aligned}$$

which defines

$$\widetilde{S}^{(2)}(s,t;u) = F_0^{-4}G^{(2)}(s,t;u) \qquad (3.28)$$

$$\widetilde{S}^{(4)}(s,t;u) = F_0^{-4}G^{(4)}(s,t;u)-\frac{1}{2}F_0^{-6}G^{(2)}(s,t;u)\sum_{i=1}^{4}(F_{P_i}^2)^{(4)}. \qquad (3.29)$$

The representation of $\phi^{(4)A}$ and $\psi^{(4)A}$ is therefore not unique. According to our definitions of the amplitude we get either

$$\phi^{(4)A}(s) = (32\pi)^2\sum_{ij}\frac{1}{z_{ij}}\frac{\overline{\overline{J}}_{ij}(s)}{F_{P_i}^2F_{P_j}^2}G_0^{(2)A_i\to ij}(s)G_0^{(2)ij\to A_f}(s)^* \qquad (3.30)$$

$$\psi^{(4)A}(s) = \frac{(96\pi)^2}{3}\sum_{ij}\frac{1}{z_{ij}}\frac{\overline{\overline{J}}_{ij}(s)}{F_{P_i}^2F_{P_j}^2}\frac{G_1^{(2)A_i\to ij}(s)G_1^{(2)ij\to A_f}(s)^*}{\lambda_{A_i}^{1/2}(s)\lambda_{A_f}^{1/2}(s)}, \qquad (3.31)$$

corresponding to the definition (3.26) or

$$\widetilde{\phi}^{(4)A}(s) = (32\pi)^2F_0^{-4}\sum_{ij}\frac{1}{z_{ij}}\overline{\overline{J}}_{ij}(s)G_0^{(2)A_i\to ij}(s)G_0^{(2)ij\to A_f}(s)^* \qquad (3.32)$$

$$\widetilde{\psi}^{(4)A}(s) = \frac{(96\pi)^2}{3}F_0^{-4}\sum_{ij}\frac{1}{z_{ij}}\overline{\overline{J}}_{ij}(s)\frac{G_1^{(2)A_i\to ij}(s)G_1^{(2)ij\to A_f}(s)^*}{\lambda_{A_i}^{1/2}(s)\lambda_{A_f}^{1/2}(s)}, \qquad (3.33)$$

when reconstructing the bare expansion of G from the "dangerous" expansion (3.27) and using the definitions (3.28, 3.29) for the $O(p^2)$ and $O(p^4)$ amplitudes.

3.3 Matching the strict chiral expansion to the dispersive representation

The dispersive representation (3.23) can be now matched to the formula (3.3). As we have mentioned above, the positions of the cuts in the formulas (3.3) and (3.23) are not the same; in the former case they correspond to the $O(p^2)$ masses (3.8), which ensures the renormalization scale independence, while in the latter they are determined by the physical ones, as required by the unitarity conditions. In order to reconcile both these requirements, one can proceed as follows (c.f. also [8]).

In (3.3), the nonanalytic terms are generally of the form $P(s)J_{ij}^r(s)$, where $J_{ij}^r(s)$ is the renormalized scalar bubble defined in Appendix D and $P(s)$ is some second order polynomial. As the first step, one rewrites these expressions in terms of $\overline{\overline{J}}_{ij}(s)$ writing $J_{ij}^r(s) = J_{ij}^r(0)+s\overline{J}_{ij}'(0)+\overline{\overline{J}}_{ij}(s)$. This adjustment allows us to split G uniquely into a polynomial part G_{pol} and a nonanalytic part G_{cut} which accumulates the unitarity cuts

$$G(s,t;u) = G_{pol}(s,t;u)+G_{cut}(s,t;u)+G\delta_G, \qquad (3.34)$$

where

$$G_{pol}(s,t;u) = \left(G(s,t;u) - G\delta_G\right)|_{\overline{J}_{ij}=\overline{\overline{J}}_{ij}=0}. \tag{3.35}$$

Both parts are now renormalization scale independent.

As a second step, we replace the $G_{cut}(s,t;u)$ with $\mathcal{G}_{unit}(s,t;u)$ from (3.23). This means we write

$$G(s,t;u) = G_{pol}(s,t;u) + \mathcal{G}_{unit}(s,t;u) + G\delta_G^{'} \tag{3.36}$$

where $\delta_G^{'}$ is a new remainder defined by this equation. According to the naive chiral power counting, $G_{cut}(s,t;u) - \mathcal{G}_{unit}(s,t;u) = O(p^6)$.

The third step, not necessary from the point of view of preserving unitarity and renormalization scale invariance, consists of a further modification of $G_{pol}(s,t;u)$ by means of replacement of the $O(p^2)$ masses $\overset{o}{M}{}_P^{2}$ in $J_{ij}^r(0)$ with the physical masses M_P^2. This replacement does not spoil the renormalization scale independence of the $G_{pol}(s,t;u)$ and corresponds to the convention introduced in [8, 9]. This again means a redefinition of the remainders $\delta_G^{'}$, *i.e.* re-shuffling of the terms of the next-to-next-to-leading order.

Note that the origin of $J_{ij}^r(0)$'s in one loop generating functional (3.5) is twofold: they can stem either from the tadpole part $Z_{tad}^{(4)}$ or from the unitarity corrections $Z_{unit}^{(4)}$. It was argued in [9] that in the former case the above mentioned replacement does not necessarily modify the numerical value of the remainders much. The reason should be that the chiral logs appear only in the combination $\mu_P \propto \overset{o}{M}{}_P^{2} \ln(\overset{o}{M}{}_P^{2}/\mu^2)$. The replacement here means

$$\overset{o}{M}{}_P^{2} \ln(\overset{o}{M}{}_P^{2}/\mu^2) \rightarrow \overset{o}{M}{}_P^{2} \ln(M_P^2/\mu^2). \tag{3.37}$$

Because $\overset{o}{M}{}_P^{2} \propto Y$, the difference should therefore either be small for $Y \sim 1$ (where $\overset{o}{M}{}_P^{2} \sim M_P^2$) or the contribution of μ_P itself is tiny for $Y \rightarrow 0$.

On the other hand, the logs from $Z_{unit}^{(4)}$ do not generally come with such a prefactor. Therefore, with a replacement $\overset{o}{M}{}_P^{2} \rightarrow M_P^2$ inside $J_{ij}^r(0)$, one might create large differences between the “old“ and “new“ remainders due to the enhancement of the contributions of chiral logs for small $Y = X/Z$. However, without the replacement inside the chiral logs of this type we could expect an unphysical increase (and irregularities) of the observables for $Y \rightarrow 0$. Also, here the replacement is natural physically, remember that the matching with the dispersive representation consist essentially of an analogous replacement within the unitary corrections. Let us also note that the splitting of the generating functional into the tadpole and unitarity part is not unique (it depends *e.g.* on the parametrization of the fluctuations around the classical solution of the $O(p^2)$ field equations in the functional integral), though the sum must be independent on this and therefore it is more consistent to use the same rule for both[12]. Nevertheless, it could be of some worth to test the differences between various treatments of the chiral logs numerically (see Subsection 6.2).

The resulting *bare* expansion (3.36) now not only meets the requirement of the exact scale independence of the remainder $\delta_G^{'}$, it has also correct physical location of the unitarity cuts. Of course, we could achieve the last property simply by inserting physical masses into the functions $\overline{\overline{J}}_{ij}(s)$ in $G_{cut}(s,t;u)$. The replacement $G_{cut}(s,t;u) \rightarrow \mathcal{G}_{unit}(s,t;u)$ has however another advantage. Namely, using the prescription (3.30), (3.31), the corresponding amplitude, written in the form

---

[12] Also notice that the offending Y dependence of the chiral logs with $O(p^2)$ masses inside comes always in the combination Y/μ^2 where μ is the renormalization scale. Provided we were able to reparametrize the bare expansion in such a way that all the running $O(p^4)$ constants were completely expressed in terms of the physical observables, the explicit independence on μ would at the same time guarantee an elimination of the irregularities for $Y \rightarrow 0$. Such a treatment has to include the reparametrization of L_1-L_3, which is, however, beyond the scope of our paper.

(without any expansion of the denominator)

$$S(s,t;u)=\frac{G(s,t;u)}{\prod_{i=1}^{4}F_{P_i}} \tag{3.38}$$

satisfies the relations of perturbative unitarity (with $S^{(2)}$ and $S^{(4)}$ given by (3.26))

$$\mathrm{disc}S_l^{(4)}(s)=\sum_{ij}\frac{2}{z_{ij}}\frac{\lambda_{ij}^{1/2}(s)}{s}S_l^{(2)ij\to A_f}(s)S_l^{(2)ij\to A_i}(s)^* \tag{3.39}$$

exactly (*i.e.* not only modulo the next-to-next-to-leading correction), which can be sometimes technically useful (*e.g.* for the unitarization by means of the inverse amplitude method, cf. [41]). The same is true using the prescription (3.32), (3.33) with $\widetilde{S}^{(2)}$ and $\widetilde{S}^{(4)}$ given by (3.28, 3.29). As we shall see in what follows, the latter prescription gives a minimal modification of the strict expansion (3.3) compatible with exact perturbative unitarity.

4. General properties of $\pi\eta$ scattering amplitude

4.1 Basic notation

Let us denote the $s-$and $u-$ channel amplitude in the isospin conservation limit as

$$\langle\pi^b(p^b)\eta(q)_{out}|\pi^a(p^a)\eta(p)_{in}\rangle=\mathrm{i}(2\pi)^4\delta(P_f-P_i)\delta^{ab}S(s,t;u) \tag{4.1}$$

and the crossed amplitude in the $t-$ channel as

$$\langle\eta(p)\eta(q)_{out}|\pi^a(p^a)\pi^b(p^b)_{in}\rangle=\mathrm{i}(2\pi)^4\delta(P_f-P_i)\delta^{ab}T(s,t;u). \tag{4.2}$$

Crossing and Bose symmetries then yield

$$\begin{aligned}T(s,t;u)&=S(t,s;u)\\ S(s,t;u)&=S(u,t;s)\\ T(s,t;u)&=T(s,u;t).\end{aligned} \tag{4.3}$$

Writing the partial wave expansion as

$$\begin{aligned}S(s,t;u)&=32\pi\sum_{l=0}^{\infty}(2l+1)P_l(\cos\theta_s)S_l(s),\\ \cos\theta_s&=\frac{(t-u)s+\Delta_{\eta\pi}^2}{\lambda_{\eta\pi}(s)},\end{aligned} \tag{4.4}$$

the scattering lengths a_l and phase shifts $\delta_l(s)$ are given by the formulae

$$\begin{aligned}\mathrm{Re}S_l(s)&=\frac{\sqrt{s}}{4}P^{2l}(a_l+O(P^2))\quad\text{for}\quad P\to 0\,,\ s\to(M_\eta+M_\pi)^2\\ \delta_l(s)&=\arctan\left(\frac{4P}{\sqrt{s}}\mathrm{Re}S_l(s)\right),\end{aligned} \tag{4.5}$$

where $P=\lambda_{\eta\pi}^{1/2}(s)/2\sqrt{s}$ is the CMS momentum. I.e., in the units of (pion Compton wavelength)$^{2l+1}$

$$a_l=M_\pi^{2l+1}\lim_{P\to 0}\frac{4}{\sqrt{s}P^{2l}}\mathrm{Re}A_l(s). \tag{4.6}$$

Let us also define the subthreshold parameters c_{ij} in terms of the expansion of the amplitude in the point of analyticity $t = 0$, $s = u = \Sigma_{\eta\pi} = M_\eta^2 + M_\pi^2$

$$S(s, t; u) = \sum_{i,j} c_{ij} t^i \nu^{2j} \tag{4.7}$$

where

$$\nu = \frac{s - u}{4M_\eta} = \frac{2s + t - 2\Sigma_{\eta\pi}}{4M_\eta}. \tag{4.8}$$

The dimension c_{ij} is $\dim[c_{ij}] = mass^{-2i-2j}$, in what follows we will refer to the dimensionless numbers $c_{ij} M_\pi^{2i+2j}$. Let us note that in the limit $m_u = m_d = 0$ we have two Adler zeros at $p^a = 0$ and $p^b = 0$, which implies the following $SU(2)_L \times SU(2)_R$ theorem

$$\lim_{m_u = m_d \to 0} c_{00} = 0. \tag{4.9}$$

We can also quote the low-energy current algebra result [42]

$$S(s, t; u) = \frac{M_\pi^2}{3F_\eta^2}, \tag{4.10}$$

which is in agreement with (4.9).

4.2 Dispersive representation

As a result of the symmetry properties of the amplitudes, the dispersive representation to the next-to-leading order (3.24) for $G_{\pi\eta}(s, t; u) = F_\pi^2 F_\eta^2 S(s, t; u)$ simplifies, namely $\phi^S = \phi^U \equiv \phi$ and $\psi^S = \psi^U \equiv \psi$. The intermediate states in (3.30, 3.31) are[13] $\pi\eta$ and $\overline{K}K$ in the s and u channels and $\pi\pi$, $\eta\eta$ and $\overline{K}K$ in the $t-$channel. This implies $\psi(s) = O(p^6)$, because the $P-$waves in the $s-$channel start at $O(p^4)$ due to the low-energy theorem (4.10) for the $\pi\eta \to \pi\eta$ amplitude and as a result of charge conjugation invariance of the $\pi\eta \to \overline{K}K$ amplitude. Moreover, $\psi^T = 0$, because the partial wave decomposition of the $t-$channel amplitude $T(s, t; u)$ contains only even partial waves due to Bose symmetry and charge conjugation. We therefore get

$$G_{\pi\eta}(s, t; u) = G_{\pi\eta,\, pol}(s, t; u) + \mathcal{G}_{\pi\eta, unit}(s, t; u) + O(p^6), \tag{4.11}$$

where the polynomial part has the following general form

$$G_{\pi\eta,\, pol}(s, t; u) = \alpha + \beta t + \gamma t^2 + \omega (s - u)^2. \tag{4.12}$$

Note that the parameters $\alpha, \ldots, \omega$ are related to the expansion of the Green function $G_{\pi\eta}(s, t; u)$ at the point of analyticity $t = 0$, $s = u = \Sigma_{\eta\pi} = M_\eta^2 + M_\pi^2$ and therefore they represent "good observables" according to our classification.

The dispersive part is

$$\mathcal{G}_{\pi\eta, unit}(s, t; u) = \phi^T(t) + \phi(s) + \phi(u), \tag{4.13}$$

where $\phi^T(t)$ and $\phi(s)$ are given by the formula (3.30). A complete list of relevant leading order contributions $G_{0,1}^{(2)12 \to ij}$ and $G_{0,1}^{(2)ij \to 34}$ can be found in Appendix C, here we give the resulting expressions (transcription to the convention (3.32, 3.33) is straightforward)

---

[13] Here we assume isospin conservation.

$$\begin{aligned}
\phi(s) \;=\; & F_0^4 \left\{ \frac{1}{9}\, \overset{o}{M}{}_{\pi}^{4}\, \frac{\overline{J}_{\pi\eta}(s)}{F_\pi^2 F_\eta^2} \right. \\
& + \frac{3}{8}[(s - \frac{1}{3}M_\eta^2 - \frac{1}{3}M_\pi^2 - \frac{2}{3}M_K^2) \\
& \left. - \frac{1}{3}(2\, \overset{o}{M}{}_{K}^{2} - \overset{o}{M}{}_{\pi}^{2} - \overset{o}{M}{}_{\eta}^{2})]^2 \frac{\overline{J}_{KK}(s)}{F_K^4} \right\}, \\
\phi^T(s) \;=\; & F_0^4 \left\{ \frac{1}{3}\, \overset{o}{M}{}_{\pi}^{2}\, [(s - \frac{4}{3}M_\pi^2) + \frac{5}{6}\, \overset{o}{M}{}_{\pi}^{2}] \frac{\overline{J}_{\pi\pi}(s)}{F_\pi^4} \right. \\
& - \frac{1}{18}\, \overset{o}{M}{}_{\pi}^{2} \left(\overset{o}{M}{}_{\pi}^{2} - 4\, \overset{o}{M}{}_{\eta}^{2} \right) \frac{\overline{J}_{\eta\eta}(s)}{F_\eta^4} \\
& + \frac{1}{8}[(s - \frac{2}{3}M_\pi^2 - \frac{2}{3}M_K^2) + \frac{2}{3}(\overset{o}{M}{}_{K}^{2} + \overset{o}{M}{}_{\pi}^{2})] \\
& \left. \times[(3s - 2M_K^2 - 2M_\eta^2) + (2\, \overset{o}{M}{}_{\eta}^{2} - \frac{2}{3}\, \overset{o}{M}{}_{K}^{2})] \frac{\overline{J}_{KK}(s)}{F_K^4} \right\}. \qquad (4.14)
\end{aligned}$$

In terms of these functions, we have (notice that $\phi^T(0) = 0$)

$$\begin{aligned}
a_0 \;=\; & \frac{1}{8\pi F_\eta^2 F_\pi^2} \frac{M_\pi}{(M_\pi + M_\eta)} \left(\alpha + 16\omega M_\eta^2 M_\pi^2 + \phi((M_\pi + M_\eta)^2) + \phi((M_\eta - M_\pi)^2)\right) \\
a_1 \;=\; & \frac{1}{12\pi F_\eta^2 F_\pi^2} \frac{M_\pi^3}{(M_\pi + M_\eta)} \left(\beta + 8\omega M_\eta M_\pi + \phi^{T'}(0) - \phi^{'}((M_\eta - M_\pi)^2)\right) \qquad (4.15)
\end{aligned}$$

and

$$\begin{aligned}
c_{00} \;=\; & \frac{1}{F_\eta^2 F_\pi^2} \left(\alpha + 2\phi(\Sigma_{\eta\pi})\right) \\
c_{10} \;=\; & \frac{1}{F_\eta^2 F_\pi^2} \left(\beta + \phi^{T'}(0) - \phi^{'}(\Sigma_{\eta\pi})\right) \\
c_{20} \;=\; & \frac{1}{F_\eta^2 F_\pi^2} \left(\gamma + \frac{1}{2}\phi^{T''}(0) + \frac{1}{4}\phi^{''}(\Sigma_{\eta\pi})\right) \\
c_{01} \;=\; & \frac{16 M_\eta^2}{F_\eta^2 F_\pi^2} \left(\omega + \frac{1}{4}\phi^{''}(\Sigma_{\eta\pi})\right). \qquad (4.16)
\end{aligned}$$

While the scattering lengths, being related to the value of the amplitude at the threshold, are not candidates for "good observables", the situation is a little bit more subtle in the case of the subthreshold parameters. Provided the η decay constant was known from experiments as accurately as F_π, then (similarly to α, β, ...) also the c_{ij} could be treated as "good observables". However, this is not the case, and we should rather use a chiral expansion of F_η in the above formulae. Therefore, the subthreshold parameters are typical examples of the dangerous ratios, which should be treated with care.

4.3 Bare expansion for $G(s,t;u)$

For the strict expansion in terms of LEC's (i.e. without any reparametrization in terms of physical observables) derived from (3.4, 3.5), we have confirmed the results of the article [38] by independent calculation. The $O(p^4)$ expansion can be written in the form

$$G_{\pi\eta} = G^{(2)} + G_{ct}^{(4)} + G_{tad}^{(4)} + G_{unit}^{(4)} + G\delta_G, \qquad (4.17)$$

where

$$
\begin{aligned}
G^{(2)}(s,t;u) &= \frac{F_0^2}{3}\overset{o}{M}{}_{\pi}^{2} \\
G^{(4)}_{ct}(s,t;u) &= 8(L_1^r(\mu)+\frac{1}{6}L_3^r(\mu))(t-2M_\pi^2)(t-2M_\eta^2) \\
&\quad +4(L_2^r(\mu)+\frac{1}{3}L_3^r(\mu))[(s-M_\pi^2-M_\eta^2)^2+(u-M_\pi^2-M_\eta^2)^2] \\
&\quad +8L_4^r(\mu)[(t-2M_\pi^2)\overset{o}{M}{}_{\eta}^{2}+(t-2M_\eta^2)\overset{o}{M}{}_{\pi}^{2}] \\
&\quad -\frac{8}{3}L_5^r(\mu)(M_\pi^2+M_\eta^2)\overset{o}{M}{}_{\pi}^{2}+8L_6^r(\mu)\overset{o}{M}{}_{\pi}^{2}(\overset{o}{M}{}_{\pi}^{2}+5\overset{o}{M}{}_{\eta}^{2}) \\
&\quad +32L_7^r(\mu)(\overset{o}{M}{}_{\pi}^{2}-\overset{o}{M}{}_{\eta}^{2})\overset{o}{M}{}_{\pi}^{2}+\frac{64}{3}L_8^r(\mu)\overset{o}{M}{}_{\pi}^{4} \\
G^{(4)}_{tad}(s,t;u) &= -\frac{F_0^2}{3}\overset{o}{M}{}_{\pi}^{2}\left(3\mu_\pi+2\mu_K+\frac{1}{3}\mu_\eta\right) \\
G^{(4)}_{unit}(s,t;u) &= \frac{1}{9}\overset{o}{M}{}_{\pi}^{4}[J^r_{\pi\eta}(s)+J^r_{\pi\eta}(u)] \\
&\quad +\frac{3}{8}[s-M_\pi^2-M_\eta^2+\frac{2}{3}\overset{o}{M}{}_{\pi}^{2}]^2J^r_{KK}(s)+\frac{3}{8}[u-M_\pi^2-M_\eta^2+\frac{2}{3}\overset{o}{M}{}_{\pi}^{2}]^2J^r_{KK}(u) \\
&\quad +\frac{1}{3}\overset{o}{M}{}_{\pi}^{2}[t-2M_\pi^2+\frac{3}{2}\overset{o}{M}{}_{\pi}^{2}]J^r_{\pi\pi}(t) \\
&\quad +\frac{2}{9}\overset{o}{M}{}_{\pi}^{2}(\overset{o}{M}{}_{\eta}^{2}-\frac{1}{4}\overset{o}{M}{}_{\pi}^{2})J^r_{\eta\eta}(t) \\
&\quad +\frac{1}{8}[t-2M_\pi^2+2\overset{o}{M}{}_{\pi}^{2}][3t-6M_\eta^2+4\overset{o}{M}{}_{\eta}^{2}-\frac{2}{3}\overset{o}{M}{}_{\pi}^{2}]J^r_{KK}(t)
\end{aligned}
\tag{4.18}
$$

are the $O(p^2)$, counterterm, tadpole and unitarity contributions respectively. In the above formulae, the masses within the loop functions $J^r_{PQ}(t)$ are the $O(p^2)$ masses

$$\overset{o}{M}{}_{\pi}^{2}=2B_0\widehat{m},\quad \overset{o}{M}{}_{K}^{2}=B_0\widehat{m}\,(r+1)\,,\quad \overset{o}{M}{}_{\eta}^{2}=\frac{2}{3}B_0\widehat{m}\,(2r+1)\,. \tag{4.19}$$

The chiral logs μ_P can be expressed using $J^r_{PP}(0)$

$$\mu_P=\frac{\overset{o}{M}{}_{P}^{2}}{32\pi^2F_0^2}\ln\frac{\overset{o}{M}{}_{P}^{2}}{\mu^2}=-\frac{\overset{o}{M}{}_{P}^{2}}{2F_0^2}\left(J^r_{PP}(0)+\frac{1}{16\pi^2}\right). \tag{4.20}$$

Written in such a form, the sum $G^{(4)}_{ct}+G^{(4)}_{tad}+G^{(4)}_{unit}$ is *exactly* renormalization scheme independent by construction. Let us now proceed as described in the previous section and write the bare expansion of $G(s,t;u)$ in the form

$$G_{\pi\eta}(s,t;u)=G_{\pi\eta,\,pol}(s,t;u)+\mathcal{G}_{\pi\eta,unit}(s,t;u)+G\delta_G^{'}. \tag{4.21}$$

Writing $J^r_{ij}(s)=J^r_{ij}(0)+\overline{J}_{ij}(s)$ in (4.18), we get the renormalization scale independent polynomial part

$$
\begin{aligned}
G_{\pi\eta,\,pol}(s,t;u) &= G^{(2)}(s,t;u)+G^{(4)}_{ct}(s,t;u)+G^{(4)}_{tad}(s,t;u) \\
&\quad +\frac{1}{3}\overset{o}{M}{}_{\pi}^{2}[t-2M_\pi^2+\frac{3}{2}\overset{o}{M}{}_{\pi}^{2}]J^r_{\pi\pi}(0) \\
&\quad +\frac{2}{9}\overset{o}{M}{}_{\pi}^{4}J^r_{\pi\eta}(0)+\frac{2}{9}\overset{o}{M}{}_{\pi}^{2}(\overset{o}{M}{}_{\eta}^{2}-\frac{1}{4}\overset{o}{M}{}_{\pi}^{2})J^r_{\eta\eta}(0)
\end{aligned}
$$

$$
\begin{aligned}
&\frac{3}{8}\{[s-M_\pi^2-M_\eta^2+\frac{2}{3}\overset{o}{M}{}_\pi^2]^2+[u-M_\pi^2-M_\eta^2+\frac{2}{3}\overset{o}{M}{}_\pi^2]^2\\
&+[t-2M_\pi^2+2\overset{o}{M}{}_\pi^2][t-2M_\eta^2+\frac{4}{3}\overset{o}{M}{}_\eta^2-\frac{2}{9}\overset{o}{M}{}_\pi^2]\}J^r_{KK}(0).
\end{aligned}
\tag{4.22}
$$

Comparing this with the general form (4.12) of $G_{\pi\eta,\,pol}(s,t;u)$, we get for the bare expansions of the parameters $\alpha-\omega$ the following manifestly renormalization scale independent form

$$
\begin{aligned}
\alpha =\; &\frac{1}{3}F_0^2\overset{o}{M}{}_\pi^2+\frac{1}{96\pi^2}\overset{o}{M}{}_\pi^2\left(\frac{7}{2}\overset{o}{M}{}_\pi^2+\frac{11}{6}\overset{o}{M}{}_\eta^2\right)\\
&+4[8(L_1^r(\mu)+\frac{1}{6}L_3^r(\mu))+\frac{3}{8}J^r_{KK}(0)]M_\pi^2M_\eta^2\\
&-[16L_4^r(\mu)+J^r_{KK}(0)]M_\pi^2\overset{o}{M}{}_\eta^2-[16L_4^r(\mu)+\frac{8}{3}L_5^r(\mu)+\frac{3}{2}J^r_{KK}(0)]M_\eta^2\overset{o}{M}{}_\pi^2\\
&-[\frac{8}{3}L_5^r(\mu)-\frac{1}{6}J^r_{KK}(0)+\frac{2}{3}J^r_{\pi\pi}(0)]M_\pi^2\overset{o}{M}{}_\pi^2\\
&+[40L_6^r(\mu)+\frac{5}{18}J^r_{\eta\eta}(0)+\frac{5}{4}J^r_{KK}(0)]\overset{o}{M}{}_\pi^2\overset{o}{M}{}_\eta^2\\
&+32L_7^r(\mu)\overset{o}{M}{}_\pi^2(\overset{o}{M}{}_\pi^2-\overset{o}{M}{}_\eta^2)\\
&+[8L_6^r(\mu)+\frac{64}{3}L_8^r(\mu)+J^r_{\pi\pi}(0)+\frac{2}{9}J^r_{\pi\eta}(0)-\frac{1}{18}J^r_{\eta\eta}(0)+\frac{1}{4}J^r_{KK}(0)]\overset{o}{M}{}_\pi^4+\frac{1}{3}F_\pi^2M_\pi^2\delta_\alpha
\end{aligned}
\tag{4.23}
$$

$$
\begin{aligned}
\beta =\; &-2\Sigma_{\eta\pi}[8(L_1^r(\mu)+\frac{1}{6}L_3^r(\mu))+\frac{3}{8}J^r_{KK}(0)]\\
&+[8L_4^r(\mu)(\overset{o}{M}{}_\eta^2+\overset{o}{M}{}_\pi^2)+\frac{1}{2}J^r_{KK}(0)\overset{o}{M}{}_\eta^2+\frac{1}{3}(J^r_{\pi\pi}(0)+\frac{1}{2}J^r_{KK}(0))\overset{o}{M}{}_\pi^2]+\beta\delta_\beta
\end{aligned}
\tag{4.24}
$$

$$
\gamma = [8(L_1^r(\mu)+\frac{1}{6}L_3^r(\mu))+\frac{3}{8}J^r_{KK}(0)]+[2(L_2^r(\mu)+\frac{1}{3}L_3^r(\mu))+\frac{3}{16}J^r_{KK}(0)]+\gamma\delta_\gamma \tag{4.25}
$$

$$
\omega = [2(L_2^r(\mu)+\frac{1}{3}L_3^r(\mu))+\frac{3}{16}J^r_{KK}(0)]+\omega\delta_\omega. \tag{4.26}
$$

5. Reparametrization of the bare expansion

Let us now discuss the various possibilities of the reparametrization of the bare expansion.

5.1 $\pi\eta$ scattering within the standard chiral perturbation theory to $O(p^4)$

The standard way of dealing with the chiral expansion consists of two "dangerous" steps. The first one involves using the inverted expansions of the type (2.10) in order to express the amplitude in terms of the masses and decay constants instead of the parameters $B_0\widehat{m}$, F_0 and $r=m_s/\widehat{m}$ of the $O(p^2)$ chiral Lagrangian. Here one encounters an ambiguity connected with different possibilities how to choose the observable G in (2.10), the chiral expansion of which starts with the desired $O(p^2)$ parameter G_0.

Let us fix this ambiguity by using the expansions of F_π^2, M_π^2 and M_K^2, inverting of which leads to[14]

$$
F_0^2 = F_\pi^2\,(1+4\mu_\pi+2\mu_K)-8M_\pi^2\,(L_4^r(\mu)(2+r)+L_5^r(\mu)) \tag{5.1}
$$

---

[14] Instead of M_K^2 we could use the chiral expansion of M_η^2 to obtain

$$
r=\widetilde{r}_2=\frac{3}{2}\left(\frac{M_\eta^2}{M_\pi^2}-\frac{1}{3}\right)
$$

$$2B_0\widehat{m} = M_\pi^2\left[1-\mu_\pi+\frac{1}{3}\mu_\eta-\frac{8M_\pi^2}{F_\pi^2}\Big(2L_8^r(\mu)+2(2+r)L_6^r(\mu)-L_5^r(\mu)-(2+r)L_4^r(\mu)\Big)\right] \quad (5.2)$$

$$r = r_2 = \frac{2M_K^2}{M_\pi^2}-1+O(p^2). \quad (5.3)$$

Inserting the inverted expansions (5.1–5.3) into (4.23) and (4.24) and keeping terms up to the order $O(p^4)$ we get

$$\begin{aligned}\alpha = {} & \frac{1}{3}F_\pi^2M_\pi^2+\frac{16}{3}M_\pi^2M_\eta^2L_3^r(\mu)-\frac{64}{3}L_7^r(\mu)M_\pi^4(r_2-1)\\ & +M_\pi^2M_\eta^2[32L_1^r(\mu)-16L_4^r(\mu)-\frac{8}{3}L_5^r(\mu)]\\ & +\frac{1}{3}(2r_2+1)M_\pi^4[32L_6^r(\mu)-16L_4^r(\mu)+\frac{2}{9}J_{\eta\eta}^r(0)]\\ & +M_\pi^4[-\frac{8}{3}L_5^r(\mu)+16L_8^r(\mu)-\frac{1}{6}J_{\pi\pi}^r(0)+\frac{2}{9}J_{\pi\eta}^r(0)-\frac{1}{18}J_{\eta\eta}^r(0)+\frac{1}{3}J_{KK}^r(0)]+\alpha\delta_\alpha^{st}\end{aligned} \quad (5.4)$$

$$\begin{aligned}\beta = {} & -2\Sigma_{\eta\pi}[8(L_1^r(\mu)+\frac{1}{6}L_3^r(\mu))+\frac{3}{8}J_{KK}^r(0)]\\ & +\frac{1}{3}M_\pi^2[16L_4^r(\mu)(r_2+2)+J_{KK}^r(0)(r_2+1)+J_{\pi\pi}^r(0)]+\beta\delta_\beta^{st},\end{aligned} \quad (5.5)$$

with new remainders δ_α^{st} and δ_β^{st}, which might be, however, out of control as we have already discussed. In fact, this first step involves three "unsafe" manipulations from the point of view of resummed χPT: using "dangerous" expansions for the masses as a starting point, the inversion and finally the negligence of all higher order terms generated by this procedure after the insertion.

It's understood to use physical masses inside the chiral logarithms. Higher order LEC's are then fitted by using additional experimental input, no parameters are therefore left free. Also note that (5.3) effectively implements the classical Gell-Mann-Okubo formula

$$3M_\eta^2-4M_K^2+M_\pi^2=0. \quad (5.6)$$

This insures renormalization scale independence. We, however, leave M_η at its physical value in cases when it was produced by on-shell mass on outer legs or inside chiral logarithms, which is compatible with the requirement of scale independence.

The second step is connected to the fact that the amplitude is used in standard χPT rather than $G(s,t;u)$. As was shown in Section 3.2, the expansion of the amplitude can be organized in various ways, of which only (3.26) is considered safe in the resummed approach. On the other hand, from the standard point of view it often seems more advantageous to use (3.28, 3.29), as together with (5.1) it leads to only the experimentally very well known pion decay constant being present in the formulae. This can be seen on the case of F_η, which is experimentally poorly known due to $\eta-\eta'$ mixing [43] and thus if it's kept at its physical value as was done in [38], a significant uncertainty is introduced into the results. As the normalization (3.28, 3.29) is used more often in NLO $S\chi PT$, we will adhere to this view and perform this second step by expanding the kaon and eta decay constants from the denominators and subsequently cutting off the higher orders.

---

or even $F_K^2M_K^2$ to get

$$r = r_2^* = 2\frac{F_K^2M_K^2}{F_\pi^2M_\pi^2}-1.$$

The latter choice, formally as good as the previous two, could also involve the redefinition of the loop masses to $\overset{o}{M}{}_P^2 = F_P^2M_P^2/F_\pi^2$ instead of the simple $\overset{o}{M}{}_P^2 = M_P^2$ as in the case of the other standard reparametrizations. Even then, however, it suffers from numerically large $O(p^4)$ corrections which could produce instabilities of the reparametrization based on this observable.

Using therefore the prescription (3.32, 3.33), the dispersive part of the $O(p^4)$ amplitude (4.14) simplifies using the reparametrization recipe described above

$$\begin{aligned}
\phi(s) &= \frac{1}{9}M_\pi^4\overline{J}_{\pi\eta}(s) \\
&+\frac{3}{8}[(s-\frac{1}{3}M_\eta^2-\frac{1}{3}M_\pi^2-\frac{2}{3}M_K^2)-\frac{1}{9}M_\pi^2(r_2-1)]^2\overline{J}_{KK}(s), \\
\phi^T(s) &= \frac{1}{3}M_\pi^2(s-\frac{1}{2}M_\pi^2)\overline{J}_{\pi\pi}(s)+\frac{1}{54}M_\pi^4\,(8r_2+1)\,\overline{J}_{\eta\eta}(s) \\
&+\frac{1}{8}[(s-\frac{2}{3}M_\pi^2-\frac{2}{3}M_K^2)+\frac{1}{3}M_\pi^2(r_2+3)] \\
&\times[(3s-2M_K^2-2M_\eta^2)+\frac{1}{3}M_\pi^2(3r_2+1)]\overline{J}_{KK}(s). \qquad (5.7)
\end{aligned}$$

The second step also propagates itself to the case of the subthreshold parameters c_{ij} and the scattering lengths a_i, where it consists of the expansion of F_η^2 in the denominator of the formulae (4.16) and (4.15). This step could in principle produce uncontrollable contribution to the remainders as well.

5.2 Resummation of the vacuum fluctuation

In order to preserve the global convergence, as was discussed, in the context of resummed χPT we are not allowed to perform "dangerous" inverted expansions and thus to express the $O(p^2)$ masses $\overset{o}{M}_P$ and the decay constant F_0 in terms of the physical ones in the way it is common within the standard χPT calculations sketched above. Instead of this, the $O(p^2)$ LEC's are left free, or more precisely, rewritten using parameters directly related to the order parameters of the chiral symmetry breaking[15]

$$r=\frac{m_s}{\widehat{m}},\quad X=\frac{\overset{o}{M}{}_\pi^2\,F_0^2}{M_\pi^2F_\pi^2},\quad Z=\frac{F_0^2}{F_\pi^2}. \qquad (5.8)$$

The bare expansions for masses $F_P^2M_P^2$ and decay constants F_P^2 are used the reparametrize the NLO LEC's L_4-L_8. As the dependence is linear, it can be done in a purely non-perturbative algebraic way by introduction of an unknown higher order remainder to each observable used. The relevant formulae for L_4-L_8 can be found in Appendix E.

As masses and decay constants do not depend on L_1, L_2, L_3, bare expansions of some additional, experimentally well known observables is needed for these LEC's. This is, however, even if highly desirable, out of the scope of our article and we make a shortcut and use the standard tabular values for these constants. We will make an analysis of the sensitivity of our results to a change in the value of L_1-L_3 in the next section devoted to numerical results.

For the resulting expression for the parameters α and β, we use the following abbreviation for some repeatedly occurring combinations

$$r_2^* = 2\frac{F_K^2M_K^2}{F_\pi^2M_\pi^2}-1 \qquad (5.9)$$

$$\varepsilon(r) = 2\frac{r_2^*-r}{r^2-1} \qquad (5.10)$$

$$\eta(r) = \frac{2}{r-1}\left(\frac{F_K^2}{F_\pi^2}-1\right) \qquad (5.11)$$

$$\Delta_{GMO} = \frac{3F_\eta^2M_\eta^2+F_\pi^2M_\pi^2-4F_K^2M_K^2}{F_\pi^2M_\pi^2}, \qquad (5.12)$$

---

[15] Here we omit the explicit dependence of X, Y and Z on N_f keeping in mind that $N_f=3$ in what follows.

in terms of which we get

$$\begin{aligned}
\alpha =\ & \frac{1}{3}XF_\pi^2M_\pi^2+\frac{1}{3}F_\pi^2M_\pi^2(1-X)\frac{5r+4}{r+2}+\frac{1}{3}F_\pi^2M_\pi^2\varepsilon(r)r\frac{2r+1}{r+2}-\frac{2}{3}\frac{F_\pi^2M_\pi^2}{r-1}\Delta_{GMO}\\
&+2\frac{F_\pi^2}{r+2}(Z-1)\left(\frac{1}{3}M_\pi^2(2r+1)+M_\eta^2\right)+\frac{F_\pi^2}{r+2}\eta(r)\left(rM_\pi^2-\frac{1}{3}(r-4)M_\eta^2\right)\\
&+\frac{1}{96\pi^2}\frac{X}{Z}M_\pi^4\,(4r+5)+\frac{3}{32\pi^2}\frac{X}{Z}M_\pi^2M_\eta^2-\frac{1}{864\pi^2}\left(\frac{X}{Z}\right)^2M_\pi^4(44r+67)\\
&-\frac{M_\pi^4}{2(r+2)(r-1)}\frac{X}{Z}[J^r_{\eta\eta}(0)(2r+1)+2J^r_{KK}(0)r-J^r_{\pi\pi}(0)(4r+1)]r\\
&+\frac{M_\pi^2M_\eta^2}{6(r+2)(r-1)}\frac{X}{Z}[J^r_{\eta\eta}(0)(2r+1)(r-4)+J^r_{\pi\pi}(0)(19r-4)-2J^r_{KK}(0)(r^2+6r-4)]\\
&+\frac{M_\pi^4}{18(r+2)(r-1)}\left(\frac{X}{Z}\right)^2[J^r_{\eta\eta}(0)(5r^2-10r-4)+6J^r_{KK}(0)(3r^2-2r-4)\\
&+4J^r_{\pi\eta}(0)(r^2+r-2)-9J^r_{\pi\pi}(0)(3r^2-2r-4)]\\
&+4[8(L_1^r(\mu)+\frac{1}{6}L_3^r(\mu))+\frac{3}{8}J^r_{KK}(0)]M_\pi^2M_\eta^2+\frac{1}{3}F_\pi^2M_\pi^2\delta_\alpha^{'},
\end{aligned} \tag{5.13}$$

$$\begin{aligned}
\beta =\ & \frac{2}{3}F_\pi^2(1-Z-\eta(r))\\
&+\frac{1}{3}\frac{M_\pi^2}{r-1}\frac{X}{Z}[J^r_{\eta\eta}(0)(2r+1)+J^r_{KK}(0)(r+1)-J^r_{\pi\pi}(0)(3r+2)-\frac{1}{16\pi^2}(r+2)(r-1)]\\
&-2\Sigma_{\eta\pi}[8(L_1^r(\mu)+\frac{1}{6}L_3^r(\mu))+\frac{3}{8}J^r_{KK}(0)]+\beta\delta_\beta^{'}.
\end{aligned} \tag{5.14}$$

The new primed remainders are the following functions of the original remainders entering the game

$$\begin{aligned}
\delta_\alpha^{'} =\ & \delta_\alpha-\frac{5r+4}{r+2}\delta_{F_\pi M_\pi}+\frac{2}{r+2}\left((2r+1)+\frac{3M_\eta^2}{M_\pi^2}\right)\delta_{F_\pi}\\
&-\frac{2}{(r+2)(r-1)}\left(3r-(r-4)\frac{M_\eta^2}{M_\pi^2}\right)(\frac{F_K^2}{F_\pi^2}\delta_{F_K}-\delta_{F_\pi})\\
&-\frac{2r(2r+1)}{(r+2)(r^2-1)}\left(2\frac{F_K^2M_K^2}{F_\pi^2M_\pi^2}\delta_{F_KM_K}-(r+1)\delta_{F_\pi M_\pi}\right)\\
&+\frac{2}{r-1}\left(3\frac{F_\eta^2M_\eta^2}{F_\pi^2M_\pi^2}\delta_{F_\eta M_\eta}+\delta_{F_\pi M_\pi}-4\frac{F_K^2M_K^2}{F_\pi^2M_\pi^2}\delta_{F_KM_K}\right)
\end{aligned} \tag{5.15}$$

$$\delta_\beta^{'} = \delta_\beta+\frac{2}{3}\frac{2F_K^2\delta_{F_K}-(r+1)F_\pi^2\delta_{F_\pi}}{\beta(r-1)}. \tag{5.16}$$

This is an alternative to the first step used in the standard approach to χPT. Because the value of F_η is not very well known, we make in a sense a parallel to the second step as well, i.e. using the chiral expansion for F_η^2 in the denominator of (4.14), (4.15) and (4.16). It involves the reparametrization in terms of X, Z, r (see Appendix B for details), but, contrary to the standard case, the denominator is not further expanded and the result is given in a nonperturbative resummed form of a ratio of two "safe" expansions.

5.3 $\pi\eta$ scattering within the generalized chiral perturbation theory to $O(p^4)$ - the bare expansion of $G(s,t;u)$

In analogy with (4.18), the strict chiral expansion for $G(s,t;u)$ within the generalized χPT can be straightforwardly obtained by using the Lagrangian summarized in the Appendix F, where we use the traditional notation for the LEC's. The result has the following structure

$$G_{\pi\eta} = \widetilde{G}^{(2)} + \widetilde{G}^{(3)} + \widetilde{G}^{(4)}_{ct} + \widetilde{G}^{(4)}_{tad} + \widetilde{G}^{(4)}_{unit} + G\delta_G^{G\chi PT}, \tag{5.17}$$

where[16]

$$\begin{aligned}
\widetilde{G}^{(2)}(s,t;u) &= \frac{1}{3}F_0^2[\widetilde{M}_\pi^2 + 4\widehat{m}^2(3A_0 - 4(r-1)Z_0^P + 2(2r+1)Z_0^S)] \\
\widetilde{G}^{(3)}(s,t;u) &= \frac{1}{3}F_0^2\Big[-2\,\widehat{m}\,\big(6\,M_\eta^2 + M_\pi^2\,(2+4\,r) - 2\,(2+r)\,t\big)\,\tilde{\xi} - 2\widehat{m}\,\Sigma_{\pi\eta}\,\xi \\
&+ 81\,\widehat{m}^3\,\rho_1 + \widehat{m}^3\,\rho_2 + \big(80 - 64\,r - 16\,r^2\big)\,\widehat{m}^3\,\rho_3 \\
&+ \big(100 + 64\,r + 34\,r^2\big)\,\widehat{m}^3\,\rho_4 + \big(2\,+\,r^2\big)\,\widehat{m}^3\,\rho_5 \\
&+ (96 - 96\,r)\,\widehat{m}^3\,\rho_6 + \big(144\,+ 288\,r + 108\,r^2\big)\,\widehat{m}^3\,\rho_7\Big]
\end{aligned}$$

$$\begin{aligned}
\widetilde{G}^{(4)}_{ct}(s,t;u) &= 8(L_1 + \frac{1}{6}L_3)(t - 2M_\pi^2)(t - 2M_\eta^2) \\
&+4(L_2 + \frac{1}{3}L_3)[(s - M_\pi^2 - M_\eta^2)^2 + (u - M_\pi^2 - M_\eta^2)^2] \\
&+\frac{8}{3}\widehat{m}^2F_0^2\,\big\{-(B_1 - B_2)\Sigma_{\pi\eta} + 2D^PM_\pi^2(r-1) - 2C_1^PM_\eta^2(r-1) \\
&+ C_1^S(2r+1)t - D^S[\frac{1}{2}\Sigma_{\pi\eta}(5r+4) - (2r+1)t] \\
&- 2B_4[3M_\eta^2 + M_\pi^2(2r^2+1) - (r^2+2)t]\big\} \\
&+\frac{1}{3}\widehat{m}^4F_0^2\,\big[256E_1 + 16E_2 + F_1^P(256 - 256r^2) + F_4^S(32 + 16r^2) \\
&+ F_1^S(256 + 320r^2) + F_5^{SP}(192 - 320r + 160r^2 - 32r^3) \\
&+ F_2^P(240 - 216r - 24r^3) + F_6^{SP}(32 - 32r + 16r^2 - 16r^3) \\
&+ F_3^P(16 - 8r - 8r^3) + F_3^S(16 + 10r + 10r^3) \\
&+ F_6^{SS}(32 + 40r + 16r^2 + 20r^3) + F_7^{SP}(384 - 160r - 256r^2 + 32r^3) \\
&+ F_2^S(400 + 234r + 74r^3) + F_5^{SS}(576 + 720r + 480r^2 + 168r^3)\big]
\end{aligned}$$

$$\begin{aligned}
\widetilde{G}^{(4)}_{tad}(s,t;u) &= -\frac{1}{9}F_0^2\,\big[2\widehat{m}B_0(\mu_\eta + 6\mu_K + 9\mu_\pi) + 8A_0\widehat{m}^2(8\mu_\eta + 3\mu_K(r+8) + 48\mu_\pi) \\
&+ 4Z_0^S\widehat{m}^2(\mu_\eta(16 + 41r) + \mu_K(48 + 90r) + \mu_\pi(96 + 45r)) \\
&- 16Z_0^P\widehat{m}^2(2\mu_\eta(5r-2) + 3\mu_K(6r-4) + 3\mu_\pi(3r-8)\big]
\end{aligned}$$

$$\begin{aligned}
\widetilde{G}^{(4)}_{unit}(s,t;u) &= \frac{1}{9}[\widetilde{M}_\pi^2 + 4\widehat{m}^2(3A_0 - 4(r-1)Z_0^P + 2(2r+1)Z_0^S)]^2[J^r_{\pi\eta}(s) + J^r_{\pi\eta}(u)] \\
&+\frac{3}{8}[s - M_\pi^2 - M_\eta^2 + \frac{2}{3}\widetilde{M}_\pi^2 - \frac{8}{3}(r-1)\widehat{m}^2(A_0 + 2Z_0^P)]^2J^r_{KK}(s) \\
&+\frac{3}{8}[u - M_\pi^2 - M_\eta^2 + \frac{2}{3}\widetilde{M}_\pi^2 - \frac{8}{3}(r-1)\widehat{m}^2(A_0 + 2Z_0^P)]^2J^r_{KK}(u) \\
&+\frac{1}{3}[\widetilde{M}_\pi^2 + 4\widehat{m}^2(3A_0 - 4(r-1)Z_0^P + 2(2r+1)Z_0^S)] \\
&\times[t - 2M_\pi^2 + \frac{3}{2}\widetilde{M}_\pi^2 + 10\widehat{m}^2(A_0 + 2Z_0^S)]J^r_{\pi\pi}(t) \\
&+\frac{2}{9}[\widetilde{M}_\pi^2 + 4\widehat{m}^2(3A_0 - 4(r-1)Z_0^P + 2(2r+1)Z_0^S)]
\end{aligned}$$

---

[16] All the LEC's in the following formulae are the renormalized LEC's at scale μ. We have omitted explicit notation of this in order to simplify the expressions.

$$
\begin{aligned}
&\times[\widetilde{M}_\eta^2-\frac{1}{4}\widetilde{M}_\pi^2+\widehat{m}^2((8r^2+1)A_0+8r(r-1)Z_0^P+2(2r+1)^2Z_0^S)]J^r_{\eta\eta}(t)\\
&+\frac{1}{8}[t-2M_\pi^2+2\widetilde{M}_\pi^2+8(r+1)\widehat{m}^2(A_0+2Z_0^S))]\\
&\times[3t-6M_\eta^2+6\widetilde{M}_\eta^2-\frac{8}{3}\widetilde{M}_K^2\\
&+\frac{8}{3}(r+1)\widehat{m}^2(3rA_0+2(r-1)Z_0^P+2(2r+1)Z_0^S]J^r_{KK}(t). \qquad (5.18)
\end{aligned}
$$

In the above formulae, the generalized $O(p^2)$ masses (also present implicitly in the chiral logs μ_P and the loop functions $J^r_{PQ}(s)$) are

$$
\begin{aligned}
\widetilde{M}_\pi^2 &= 2[B_0+2\widehat{m}(r+2)Z_0^S]\widehat{m}+4A_0\widehat{m}^2\\
\widetilde{M}_K^2 &= [B_0+2\widehat{m}(r+2)Z_0^S]\widehat{m}\,(r+1)+A_0\widehat{m}^2\,(r+1)^2\\
\widetilde{M}_\eta^2 &= \frac{2}{3}[B_0+2\widehat{m}(r+2)Z_0^S]\widehat{m}\,(2r+1)+\frac{4}{3}A_0\widehat{m}^2\left(2r^2+1\right)+\frac{8}{3}Z_0^P\widehat{m}^2(r-1)^2. \qquad (5.19)
\end{aligned}
$$

The unitarity part can be further split into the polynomial and dispersive part

$$
\widetilde{G}^{(4)}_{unit}=\widetilde{G}^{(4)}_{unit,\,pol}+\widetilde{G}^{(4)}_{unit,\,disp}=\widetilde{G}^{(4)}_{unit}|_{J^r\to J(0)}+\widetilde{G}^{(4)}_{unit}|_{J^r\to\bar{J}}. \qquad (5.20)
$$

According to the general prescription, the dispersive part can be replaced with that of the dispersive representation which has the general form

$$
\widetilde{\mathcal{G}}^{(4)}_{unit}=\phi^T(t)+\phi(s)+\phi(u), \qquad (5.21)
$$

where now

$$
\begin{aligned}
\phi(s) &= F_0^4\left\{\left(\frac{1}{3}\alpha_{\pi\eta}\widetilde{M}_\pi^2\right)^2\frac{\overline{J}_{\pi\eta}(s)}{F_\pi^2F_\eta^2}\right.\\
&\left.+\frac{3}{8}[s-\frac{1}{3}M_\eta^2-\frac{1}{3}M_\pi^2-\frac{2}{3}M_K^2-\frac{1}{3}(2\widetilde{M}_K^2-\widetilde{M}_\pi^2-\widetilde{M}_\eta^2+\alpha_{\pi\eta K}\widetilde{M}_\pi^2)]^2\frac{\overline{J}_{KK}(s)}{F_K^4}\right\}
\end{aligned}
\qquad (5.22)
$$

$$
\begin{aligned}
\phi^T(s) &= F_0^4\left\{\frac{1}{3}\alpha_{\pi\eta}\widetilde{M}_\pi^2[s-\frac{4}{3}M_\pi^2+\frac{5}{6}\alpha_{\pi\pi}\widetilde{M}_\pi^2]\frac{\overline{J}_{\pi\pi}(s)}{F_\pi^4}\right.\\
&-\frac{1}{18}\alpha_{\eta\eta}\alpha_{\pi\eta}\widetilde{M}_\pi^2\left(\widetilde{M}_\pi^2-4\widetilde{M}_\eta^2\right)\frac{\overline{J}_{\eta\eta}(s)}{F_\eta^4}\\
&+\frac{1}{8}[s-\frac{2}{3}M_\pi^2-\frac{2}{3}M_K^2+\frac{2}{3}((\widetilde{M}_K-\widetilde{M}_\pi)^2+2\alpha_{\pi K}\widetilde{M}_K\widetilde{M}_\pi]\\
&\left.\times[3s-2M_K^2-2M_\eta^2+\alpha_{\eta K}(2\widetilde{M}_\eta^2-\frac{2}{3}\widetilde{M}_K^2)]\frac{\overline{J}_{KK}(s)}{F_K^4}\right\}
\end{aligned}
\qquad (5.23)
$$

for (3.30, 3.31) and analogously for (3.32, 3.33). The coefficients $\alpha_{\pi\eta}\ldots$ parametrize the difference between the standard and the generalized cases, within the standard $O(p^4)$ chiral expansion their values are either one or zero. The dependence of these constants on the LEC's are given in the Appendix G.

5.4 Observables of the $\pi\eta$ scattering within $G\chi PT$ to $O(p^4)$ - the reparametrization

As it can be easily seen from the above formulae (in fact it is a consequence of the construction of

$G\chi PT$), after identifying the parameters of the Lagrangians,

$$\begin{aligned}
\frac{B_0\widehat{m}}{F_0^2}L_4^r(\mu) &\rightarrow \frac{1}{8}\widehat{m}\widetilde{\xi} \\
\frac{B_0\widehat{m}}{F_0^2}L_5^r(\mu) &\rightarrow \frac{1}{8}\widehat{m}\xi^r \\
\frac{B_0^2\widehat{m}^2}{F_0^2}L_6^r(\mu) &\rightarrow \frac{1}{16}\widehat{m}^2 Z_0^S \\
\frac{B_0^2\widehat{m}^2}{F_0^2}L_7^r(\mu) &\rightarrow \frac{1}{16}\widehat{m}^2 Z_0^P \\
\frac{B_0^2\widehat{m}^2}{F_0^2}L_8^r(\mu) &\rightarrow \frac{1}{16}\widehat{m}^2 A_0
\end{aligned} \tag{5.24}$$

and defining the remainders using the *physical* masses inside the chiral logarithms and the loop functions J_{PQ}^r, the generalized *bare* chiral expansions contain all the terms of the standard one. More precisely, the generalized $O(p^4)$ bare expansions include extra $O(p^4)$ terms, which are counted as $O(p^6)$ and $O(p^8)$ within the standard chiral power counting scheme. As a consequence, after writing the generalized bare chiral expansion of a generic "good" observable g in the form

$$g = g^{(2),\,G\chi PT} + g^{(3),\,G\chi PT} + g^{(4),\,G\chi PT} + g\delta_g^{G\chi PT} \tag{5.25}$$

and then collecting the "standard" terms together, this expansion can be formally rewritten as

$$g = g^{(2),\,std} + g^{(4),\,std} + g\delta_g, \quad g\delta_g = g\delta_g^{(G)} + g\delta_g^{G\chi PT}, \tag{5.26}$$

where the identification (5.24) is assumed. The extra $O(p^4)$ terms mentioned above are now accumulated in $g\delta_g^{(G)}$. In the case of the polynomial parameters $\alpha \ldots \omega$ (4.23–4.26), the two versions of the chiral expansion coincide for γ and ω[17]

$$\delta_\gamma = \delta_\gamma^{G\chi PT} \qquad \delta_\omega = \delta_\omega^{G\chi PT}, \tag{5.27}$$

while the "standard" remainders δ_α, δ_β can be split into an explicitly known part, which includes the extra "nonstandard" terms, and the unknown remainders inherent to $G\chi PT$

$$\delta_\alpha = \delta_\alpha{}^{loops}(\mu) + 3\frac{\widehat{m}^2F_0^2}{F_\pi^2M_\pi^2}\delta_\alpha^{CT}(\mu) + \delta_\alpha^{G\chi PT} \tag{5.28}$$

$$\beta\delta_\beta = \beta\delta_\beta{}^{loops}(\mu) + \widehat{m}^2F_0^2\delta_\beta^{CT}(\mu) + \beta\delta_\beta^{G\chi PT}. \tag{5.29}$$

Here the first terms correspond to the new loops and the second to the new counterterm contributions. The explicit expressions for them can be easily extracted from the formulae of the previous subsection, the results are however rather lengthy and we postpone them to Appendix H.

Let us note that both $\delta_{\alpha,\beta}^{loop}(\mu)$ and $\delta_{\alpha,\beta}^{CT}(\mu)$ are generally renormalization scale dependent. However, due to the running of the $G\chi PT$ LEC's A_0, Z_0^S, Z_0^P, ξ and $\widetilde{\xi}$,which, after identification (5.24), is the same as in the standard case, the "standard" remainders δ_α and δ_β given by (5.28, 5.29) are μ-independent. Of course, the "true $G\chi PT$" remainders $\delta_\alpha^{G\chi PT}, \ldots, \delta_\omega^{G\chi PT}$ are scale independent by construction. That means the sum of the loop and counterterm contributions to the "standard" remainders is μ-independent too.

The usual way how to handle the reparametrization of the $G\chi PT$ bare expansions is quite similar to the standard one. The difference is that as there are three additional $O(p^2)$ LEC's, not all of them can be reparametrized using inverted mass and decay constant expansions. The solution is to leave two of them free (*e.g.* r and $\zeta = Z_0^S/A_0$). Consequently, the expansion is performed according to the generalized power counting scheme and the terms of order higher than $O(p^4)$ are discarded.

---

[17] The reason is that they stem from the terms quadratic in the Mandelstam variables.

We will, however, not use this approach, but rather exploit the relation (5.26), *i.e.* sew the standard and generalized bare expansions together. The reparametrization is then an extension of the resummed one (Appendix E), where all the remainders of mass and decay constant bare expansions are split according to (5.26). We can use all the resummed formulae as they are exact algebraic identities, valid independently on the version of χPT. The generalized contributions to the remainders can be found in Appendices B and I. The outcome for the parameters α and β is then obtained by simply inserting all the generalized results for the remainders (Appendices B, H, I) into the expression for $\delta_\alpha^{'}$ and $\delta_\beta^{'}$ (5.15, 5.16).

After this procedure, the generalized LEC's are present only in the formulae for the standard remainders. Also note that ${\delta_\alpha}^{loops}(\mu)$, ${\delta_\beta}^{loops}(\mu)$ as well as the generalized loop contributions to mass and decay constant remainders depend explicitly on the $O(p^2)$ LEC's $B_0 = XM_\pi^2/2\widehat{m}$, $F_0^2 = ZF_\pi^2$, A_0, Z_0^S and Z_0^P.[18] So as the last step of the reparametrization, the remaining dependence of the generic "loop" remainders ${\delta_\alpha}^{loops}(\mu), \ldots, etc.$ on the $O(p^2)$ LEC's F_0, A_0, Z_0^S and Z_0^P can be removed up to the order $O(p^4)$ using the leading order expressions

$$\begin{aligned}
F_0^2 = F_K^2 = F_\eta^2 &\rightarrow F_\pi^2 \\
\widehat{m}^2 F_0^2 Z_0^S &\rightarrow \frac{1}{4}\frac{F_\pi^2 M_\pi^2}{r+2}(1 - X - \varepsilon(r)) \\
\widehat{m}^2 F_0^2 Z_0^P &\rightarrow -\frac{1}{8}F_\pi^2 M_\pi^2\left(\varepsilon(r) - \frac{\Delta_{GMO}}{(r-1)^2}\right) \\
\widehat{m}^2 F_0^2 A_0 &\rightarrow \frac{1}{4}F_\pi^2 M_\pi^2 \varepsilon(r).
\end{aligned} \tag{5.30}$$

As a summery, our handling of the generalized bare expansion can be viewed in two ways - either as a partial estimate of the standard remainders present in the resummed approach or as a special treatment within the generalized framework, where the $O(p^2)$ (and partly $O(p^3)$) LEC's are reparametrized algebraically at the leading order, while treated perturbatively at the $O(p^4)$ one. The numerical results including a simple estimate of the remaining NLO and NNLO LEC's are presented in Subsection 6.6, also Appendix B contains an illustrative example of applying this procedure on F_η.

6. Numerical results

In this section we shall present the numerical analysis of the observables connected to the $\pi\eta-$ scattering amplitude and the results which illustrate the subtleties of the various versions of the chiral expansions described above. In the numerical estimates we use $M_\pi = 135\mathrm{MeV}$, $M_\eta = 548\mathrm{MeV}$, $M_K = 496\mathrm{MeV}$, $\mu = M_\rho = 770\mathrm{MeV}$, $F_\pi = 92.4\mathrm{MeV}$ and $F_K = 113\mathrm{MeV}$. For the calculation within the standard χPT, the $O(p^4)$ LEC's are taken from ref. [44] and [45]. In the alternative reparametrization schemes, where only L_1, L_2 and L_3 remain among the free parameters and the other $O(p^4)$ LEC's are expressed in terms of physical masses, decay constants and the indirect remainders, we again keep (though rather non-systematically) the values of L_1, L_2 and L_3 from the same references. The sensitivity on this LEC's might be then estimated by means of the variation around these central values. In this chapter we insert the physical masses into the functions $J_{PQ}^r(0)$ unless stated otherwise.

---

[18] More precisely, the loops depend on the "true $O(p^2)$ LEC's"$\overline{A}_0$, $\overline{Z}_0^{S,P}$ (cf. Appendix F), the difference is however of the higher order in the generalized power counting.

6.1 The standard chiral perturbation theory

L_i	α/α^{CA}	$10^3\beta/M_\eta^2$	$10^3\gamma$	$10^4\omega$
[44]	1.68	0.90	−1.52	2.24
[45]	1.91	−0.68	−0.23	−5.03
Δ	2.48	7.49	3.31	9.48

Table 1: The standard $O(p^4)$ values of the polynomial parameters for the two sets of LEC's taken form references [44] and [45]. In the last row, the sensitivity on the LEC's is (over)estimated adding the uncertainties associated with the LEC's [44] in quadrature (This is of course only a rough estimate, because in fact not all the uncertainties of the L_i's are independent).

This subsection discusses the predictions of the standard chiral expansion to the order $O(p^4)$, which are summarized in Table 1 and 2. Let us start with the parameter α of the polynomial part of the amplitude. The relevant formulae from the Subsection 5.1 and the LEC's taken from [44][19] result in the following value

$$\alpha = \frac{1}{3}\left(1 + 0.683 + \delta_\alpha^{st}\ \right) F_\pi^2 M_\pi^2 \tag{6.1}$$

In this expression, the first term corresponds to the current algebra result $\alpha^{CA} = F_\pi^2 M_\pi^2/3$, while the second one represents the $O(p^4)$ correction. The third term is the standard remainder, which might be out of control when $X, Z \ll 1$ and r far from r_2, even if the bare expansion of α were globally convergent (let us remind that α is a "good" observable) as we have discussed in Section 2. Let us also notice the unusually large next-to-leading correction, which could also indirectly indicate the numerical importance of the remainder in this scheme.

The actual numerical value of the NLO correction is very sensitive to a shift in the $O(p^4)$ LEC's. The corresponding variation $\Delta\alpha$ is numerically

$$\frac{\Delta\alpha}{\alpha^{CA}} = (3.38\Delta L_1 + 0.56\Delta L_3 - 3.50\Delta L_4 - 0.30\Delta L_5 + 3.62\Delta L_6 - 3.42\Delta L_7 + 0.10\Delta L_8) \times 10^3. \tag{6.2}$$

For example, using the $O(p^6)$ analysis based LEC's from [45] instead of those from [44], we get (cf. Table 1)[20]

$$\frac{\Delta\alpha}{\alpha^{CA}} = 0.23. \tag{6.3}$$

Note that the large coefficients in front of the L_4 and L_6 contributions indicate sensitivity of this observable to the vacuum fluctuations of $\overline{s}s$ pairs as mentioned in the Introduction.

L_i	c_{00}/c_{00}^{CA}	10^3c_{10}	10^3c_{20}	10^3c_{01}	a_0/a_0^{CA}	10^3a_1
[44]	1.06	0.91	−1.23	8.27	1.96	0.59
[45]	1.51	−0.67	0.07	−3.36	1.18	−0.60
Δ	2.49	7.49	3.31	15.16	3.21	2.80

Table 2: The standard $O(p^4)$ values of the subthreshold and threshold parameters as in Tab. 1. The c_{ij} parameters are given in their natural units described in the main text.

Let us compare this case with the related "dangerous" observable, namely the subthreshold parameter c_{00}. From (4.16) we get

$$c_{00} = \frac{1}{3}\left(1 + 0.683 - 0.625 + 0.006\right)\frac{M_\pi^2}{F_\pi^2} = 1.064 c_{00}^{CA}, \tag{6.4}$$

where the individual terms are the leading order contribution $c_{00}^{CA} = M_\pi^2/3F_\pi^2$, the next-to-leading correction to the parameter α, the next-to-leading correction to F_η^2 induced by the expansion of the denominator and the contribution stemming from the unitarity correction $\phi(s)$ respectively. The first two large corrections accidentally cancel here, this, however, does not automatically imply similar cancellation of the potentially large remainders (we have not written them down explicitly here). Also, the strong sensitivity of α to the variation of the LEC's propagates here giving

---

[19] This set of LEC's is used in numerical estimates unless stated otherwise.

[20] Because the values of the LEC's L_i based on the $O(p^6)$ fit include implicitly parts of the $O(p^6)$ corrections, the large variation can be interpreted as a signal of the importance of the NNLO contributions to the parameter α. The same is true for other observables from the Table 1.

$$\frac{\Delta c_{00}}{c_{00}^{CA}} = \frac{\Delta\alpha}{\alpha^{CA}} - 0.28\Delta L_5 \times 10^3 \tag{6.5}$$

and furthermore increases the uncertainty of the $O(p^4)$ correction. This strong sensitivity supports the possibility that the standard remainders for c_{00} might be numerically larger than the next-to-leading correction. Namely using the LEC's from the $O(p^6)$ fit [45], which generates part of the $O(p^6)$ corrections to the reparametrized expansion of c_{00}, we get

$$\frac{\Delta c_{00}}{c_{00}^{CA}} = 0.45. \tag{6.6}$$

We can also check the sensitivity of the next-to-leading order contributions to the way we rewrite them in terms of the physical masses and decay constants (*i.e.* how we use the $O(p^2)$ relations generating here a difference of the order $O(p^6)$). Provided we insert the alternative $O(p^2)$ expressions for r into the chiral expansions of α and c_{00}

$$\widetilde{r}_2 = \frac{1}{2}\left(\frac{3M_\eta^2}{M_\pi^2} - 1\right) = 24.2 \tag{6.7}$$

$$r_2^* = 2\frac{F_K^2 M_K^2}{F_\pi^2 M_\pi^2} - 1 = 39.4, \tag{6.8}$$

instead of the standard $O(p^2)$ value $r = r_2 = 2M_K^2/M_\pi^2 - 1 = 25.9$, we get as a result

$$\widetilde{\alpha} = \frac{1}{3}(1 + 0.601)\, F_\pi^2 M_\pi^2 \tag{6.9}$$

$$\widetilde{c}_{00} = \frac{1}{3}(1 + 0.031)\frac{M_\pi^2}{F_\pi^2} \tag{6.10}$$

and

$$\alpha^* = \frac{1}{3}(1 + 1.297)\, F_\pi^2 M_\pi^2 \tag{6.11}$$

$$c_{00}^* = \frac{1}{3}(1 + 0.325)\frac{M_\pi^2}{F_\pi^2}. \tag{6.12}$$

The remaining parameters of the polynomial part start at $O(p^4)$ and we get them from (5.5), (4.25) and (4.26). Their numerical values and the related subthreshold parameters c_{ij} in natural units (chosen in such a way to make the comparison with the polynomial parameters easy, *i.e.* we take c_{10} and c_{01} in units of M_η^2/F_π^2 and c_{20} in units of F_π^{-4}, cf. (4.16)) are shown in Tables 1 and 2 for the two sets of $O(p^4)$ LEC's.

All the considered parameters are strongly sensitive to the variations of the LEC's. For instance the parameter β varies with the L_i's as

$$\Delta\beta = (17.0\Delta L_1 - 2.8\Delta L_3 + 9.1\Delta L_4)M_\eta^2. \tag{6.13}$$

For the LEC's [44] we get $\beta = 0.90 \times 10^{-3} M_\eta^2$. Using the set [45] we get a drastic change

$$\Delta\beta = -1.58 \times 10^{-3} M_\eta^2. \tag{6.14}$$

Let us turn to the "doubly dangerous" observables represented by the scattering lengths now. For the s-wave we obtain from (4.15) and the LEC's [44]

$$\begin{aligned} a_0 &= \frac{1}{24\pi F_\pi^2}\frac{M_\pi^3}{M_\eta + M_\pi}(1 + 0.683 + 0.378 - 0.625 + 0.527) \\ &= \frac{1}{24\pi F_\pi^2}\frac{M_\pi^3}{M_\eta + M_\pi}(1 + 0.963) = 11.0 \times 10^{-3}. \end{aligned} \tag{6.15}$$

Here the individual terms in the first line represent the current algebra result, the correction stemming from the $O(p^4)$ contributions to the parameters α and ω, the next-to-leading correction to F_η^2 induced by the expansion of the denominator and the correction induced by the dispersive part of the amplitude $\phi(s)$, in this order. This result confirms the expectations about a bad convergence of the chiral expansion for the observables which are connected to the threshold values of the amplitude - even if the polynomial NLO corrections were small, which are not, the dispersive part would be still as large as 50% of the leading order term.

The sensitivity to the $O(p^4)$ LEC's is illustrated in Table 2. The p-wave scattering length then starts at $O(p^4)$, we get the values in the last column of the table from (4.15).

When comparing our results for the scattering lengths with those of [38], quoted in the Introduction, we can see a large discrepancy

$$a_0^{BKM} = 7.2 \times 10^{-3} \qquad a_1^{BKM} = -5.2 \times 10^{-4}. \tag{6.16}$$

The difference is produced by a different set of $O(p^4)$ LEC's, the alternative treatment of the F_η in the denominator and by another form of the unitarity corrections - the authors do not use a matching with a dispersive representation. Taken these distinctions into account, we get more consistent numbers (with our inputs for the masses and decay constants): $a_0 = 7.0 \times 10^{-3}$ and $a_1 = -5.0 \times 10^{-4}$. As we can see, a slightly different treatment of the standard chiral expansion may lead to a significant shift in results.

6.2 Resummation of vacuum fluctuations - basic properties

In the resummed case, the free parameters are X, Z, and r together with the remaining LEC's L_1, L_2 and L_3 and the direct and indirect remainders $\delta_\alpha \ldots \delta_\omega$ and δ_{F_P}, $\delta_{F_P M_P}$. Because F_η is experimentally not known with enough accuracy, we also have to fix how to treat the observable Δ_{GMO} which was introduced to eliminate the LEC L_7 using the bare expansion for $F_\eta^2 M_\eta^2$. Let us remind that our definition of Δ_{GMO} follows the ref. [8], where it is based on the good observables $F_P^2 M_P^2$ instead of M_P^2 and differs from that originally defined in [3]. One possibility is to treat Δ_{GMO} as an additional independent parameter. The another, similarly to the treatment of F_η in the denominators of (4.14), (4.15) and (4.16) discussed earlier, is to use a (resummed) chiral expansion of F_η^2 inserted to Δ_{GMO} for the numerical estimates, *i.e.* to insert the following exact algebraic identity into (5.13) (cf. Appendix B for details)

$$\begin{aligned} F_\pi^2 M_\pi^2 \Delta_{GMO} &= F_\pi^2 M_\pi^2 - 4M_K^2 F_K^2 + M_\eta^2 (1-\delta_{F_\eta})^{-1} \Bigg(4F_K^2(1-\delta_{F_K}) - F_\pi^2(1-\delta_{F_\pi}) \\ &\quad - M_\pi^2 \left(\frac{X}{Z}\right) \left(J_{\pi\pi}^r(0) - 2(r+1)J_{KK}^r(0) + (2r+1)J_{\eta\eta}^r(0)\right)\Bigg). \end{aligned} \tag{6.17}$$

This generates the indirect remainder δ_{F_η} in a nonlinear way.

Before doing a more detailed analysis, let us first illustrate the numerical sensitivity connected with the subtleties of the definition of the bare expansion. As we have discussed in Section 3, there is still some freedom how to define the amplitudes entering the dispersive part of the $G_{\pi\eta}$ (cf. (3.30, 3.31) and (3.32, 3.33) and also how to treat the masses inside the chiral logs. Based on general considerations it was argued [9], that in the latter case the different prescriptions should not make much difference. Nevertheless, it might be interesting to test this assumption numerically in our concrete case and also to check what is the numerical influence of the varying amplitude definition.



Figure 1: *Comparison of the numerical impact of the various forms of the dispersive part on the scattering length a_0. The full line represents the strict chiral expansion, dotted, dashed and dash-dotted lines the "minimal" modification, (3.30, 3.31) and (3.32, 3.33) respectively. The horizontal line shows our standard NLO prediction.*

In Figure 1 we plot the comparison of various definitions of the dispersive part of the amplitude using the scattering length a_0 as an example, *i.e* we illustrate its sensitivity on the various versions of the unitarity corrections. The cusps on the full line, which uses the *strict* chiral expansion with the unphysical choice of the $O(p^2)$ masses in all J^r_{ij}, originate in the conflict of the physical masses used for the on-shell outer legs and the unphysical location of the thresholds. This illustrates the fact that the original strict chiral expansion is unsuitable for realistic physical predictions and its redefinition into a bare one is necessary. The dotted line shows the "minimal" physical modification of the strict expansion by means of insertion physical masses into all J^r_{ij}. While the "minimal" version and the unitary choice (3.32, 3.33) give numerically almost the same result, the difference between the these two and the third possibility (3.30, 3.31) is up to $\sim 0.3\ a_0^{CA}$.



Figure 2: *In this figure we illustrate the sensitivity of the "good" variables α and β to the treating of the chiral logs. The full line corresponds to the $O(p^2)$ masses in all $J^r_{ij}(0)$'s, while doted and dashed lines to the physical masses either in all $J^r_{ij}(0)$'s or only in $J^r_{ij}(0)$'s originating from the unitarity corrections. Horizontal lines are standard NLO predictions.*

Figure 2 shows the dependence of the polynomial parameters α and β on $Y = X/Z$ for $Z = 0.8$ and $r = 20$, using the the various possible treatments of chiral logarithms in the bare expansion. The results demonstrate that the difference might be numerically important in some range of Y. For α the various possibilities do not differ drastically in comparison with the value of α itself, on the other hand the differences become comparable with α^{CA} at $Y \sim 0.5$. As we have discussed in Section 2, in the region of small Y the case with $O(p^2)$ masses in the tadpoles only should not differ drastically from the case when all the masses are physical. However, the convergence to the common value at $Y = 0$ is rather slow and in the intermediate region of Y the difference of this two cases for α is $\sim 0.5\ \alpha^{CA}$ in a relatively wide interval. Keeping $O(p^2)$ masses also in the unitarity corrections produce instabilities for $Y \to 0$ as expected. The parameter β (which starts at $O(p^4)$) is even much more sensitive.

In the following numerical analysis we take a pragmatic point of view and fix the bare expansion in such a way that the comparison of the resummed and standard reparametrizations remains as simple as possible, *i.e.* we insert physical masses into $J^r_{ij}(0)$ and define the amplitude according to (3.28, 3.29) and (3.32, 3.33).

6.3 Numerical comparison of the resummed and standard reparametrization

Within the standard χPT we have an $O(p^4)$ prediction for X, Z, r and Δ_{GMO} based on the standard formal $O(p^4)$ chiral expansion (A.1) and (A.3), cf. Appendix A. Using the LEC's from [44] and [45], we get numerically the following central values, which should confirm the self-consistency of the standard chiral expansion scheme

L_i set	X^{std}	Z^{std}	r^{std}	r^{*std}	Δ^{std}_{GMO}
[44]	0.902	0.865	25.2	26.7	6.41
[45]	0.726	0.734	25.9	31.7	3.31

As we can see, while expectations are fulfilled in the first case, there is a considerable shift when using the $O(p^6)$ fitted constants. These numbers, moreover, should be taken with some caution, because they originate in the expansions of the "dangerous" observables and can be therefore plagued with large $O(p^6)$ remainders as well as with strong sensitivity to the $O(p^4)$ LEC's[21]. In the above table r^{std} stems from the chiral expansion of r_2 while r^{*std} uses expansion of r_2^*. [22]

X	Z	r	Δ_{GMO}	α/α^{CA}	$10^3\beta$	c_{00}/c_{00}^{CA}	$10^3 c_{10}$	a_0/a_0^{CA}	$10^3 a_1$
X^{std}	Z^{std}	r^{std}	Δ^{std}_{GMO}	1.88	0.69	1.11	0.41	1.64	0.30
X^{std}	Z^{std}	r^{*std}	Δ^{std}_{GMO}	1.61	0.55	0.95	0.33	1.47	0.26
X^{std}	Z^{std}	r_2	Δ^{std}_{GMO}	1.74	0.62	1.03	0.37	1.55	0.28
Z^{std}	Z^{std}	r^{*std}	Δ^{std}_{GMO}	1.76	0.75	1.04	0.45	1.57	0.31
Z^{std}	Z^{std}	r^{std}	Δ^{std}_{GMO}	2.02	0.89	1.20	0.53	1.72	0.35
X^{std}	Z^{std}	r^{std}	Δ_{GMO}	2.07	0.69	1.22	0.41	1.75	0.30
X^{std}	Z^{std}	r^{*std}	Δ_{GMO}	1.78	0.55	1.05	0.33	1.57	0.26

Table 3: *The values of the the polynomial parameters α and β and the related subthreshold and threshold parameters near the standard reference point. For Δ_{GMO} we take either the standard value (Δ^{std}_{GMO}) or the resummed prediction described (Δ_{GMO}) in the main text.*

Let us now illustrate the relationship of the resummed and standard approach using the observables from Subsection 6.1.

For the "good" observables α and β we can expect that the numerical values of X^{std}, Z^{std}, r^{std} and Δ^{std}_{GMO}, with L_1, L_2 and L_3 taken from [44] for definiteness, should produce numbers consistent with the first row of Table 1 when inserted into (5.13, 5.14). The results for the various possibilities how to approach the standard predictions for α and β (which is independent on Δ_{GMO}) within

---

[21] As it was analyzed in detail in [8], the actual values of X^{std} and Z^{std} are strongly sensitive to the values of the LEC's L_6 and L_4 connected with the vacuum fluctuation of the $\bar{s}s$ pairs, the same is true for the sensitivity of r^{std} and Δ^{std}_{GMO} to L_8 and L_7. This causes large error bars to be attached to these values. Nevertheless, in the following we take these central values as a reference point for an illustrative numerical comparison of the two versions of the chiral expansion.

[22] Though the difference between the values of r^{std} and r^{*std} is within the standardly expected accuracy of the $O(p^4)$ approximation, note, however, that for r^{*std} the $O(p^4)$ correction is much larger than in the first alternative ($r_2 = 25.9$ while $r_2^* = 39.4$).



Figure 3: *The dependence of the parameters α and β on r, X and Z is plotted, one of the parameters being fixed at its standard reference value in each figure. The dashed horizontal line shows the standard values from the first row of the Table 1, the full circle depicts the corresponding resummed value at the standard reference point $[r^{std}, X^{std}, Z^{std}]$. The error bars represent the* 10% *uncertainties from the direct and indirect remainders added in quadrature. In the first row, r is fixed at r^{std}, the filled are highlight the dependence on Z between $Z = Z^{std}$ (solid line) and $Z = 0.5$ (dotted one). Similarly, in the second row $X = X^{std}$, the filled area shows the dependence on Z again. Z is fixed at Z^{std} in the last row, the solid line shows the case with $r = r^{std}$ while the dotted the one with a lower value $r = 15$.*

the resummed version of χPT are summarized[23] in the Table 3. The last row corresponds to the resummed treating of Δ_{GMO} explained above. The dependence of the central values of α and β on the parameters r, X and Z in the broader vicinity of their standard values is illustrated in Fig.3. These results can be interpreted as a consistency of both variants of reparametrization for good observables near the standard reference point X^{std}, Z^{std} and r^{std}, where the predictions of the resummed version almost coincide with the standard results[24]. This coincidence together with the working hypothesis about the controllable remainders of good observables within the resummed

---

[23] In this and the following tables in this subsection we ignore the uncertainty stemming from the remainders and L_i, $i = 1, 2, 3$ and give only the central values (assuming the central values of the remainders to be zero).

[24] As a rule, the point X^{std}, Z^{std} and r^{std} cannot give the best coincidence with the standard values in all cases,



Figure 4: *Dependence of the subthreshold parameters* c_{00} *and* c_{10} *related to the polynomial parameters* α *and* β*. The figures are in one-to one correspondence to those in Figure 3.*

reparametrization scheme confirms again a self-consistency of the standard expansion based on the assumption $X \sim 1$, $Z \sim 1$ and $r \sim r_2$. Away from the standard reference point, however, the standard reparametrization might be dangerous in the sense that the difference between the standard and the resummed prediction diverges rapidly and the importance of the standard $O(p^6)$ remainders might therefore increase considerably.

For the "dangerous" observables like c_{ij} we cannot *a priori* expect coincidence of both expansion even near the standard values of X, Z, and r due to the different treatment of the denominators, which contain large $O(p^4)$ corrections and are not expanded in the resummed case. Comparison of both approaches is illustrated in Table 3 (with the same treatments of Δ_{GMO} as above), Table 4 and in Fig. 4. For the dispersive part we use the prescription (3.32, 3.33), which differs from the corresponding standard contributions of the unitarity corrections to c_{ij} for $X = Z = 1$ and $r = r_2$ by a factor $F_\pi^2/F_\eta^2 \approx 0.6$. This is reflected by the values of those c_{ij} that start at $O(p^4)$ (cf. Table 3, 4). Namely in this case the contribution of the polynomial part is reduced near the reference point roughly by the same factor with respect to the standard value (which includes only the first term of the expansion of the denominator). On the other hand, c_{00} is compatible with the standard

---

the reason can be understood *e.g.* by having a closer look on the resummed reparametrization of β (cf. (5.14)). In order to reproduce the dependence of β on L_4 satisfactorily, we need $Z = Z^{std}$ and $r = r^{std}$, on the other hand to reproduce the chiral logs we need rather $X/Z = 1$ and $r = r_2$. This can explain why β approaches the standard value best for $X = Z = Z^{std}$.



Figure 5: *Dependence of the scattering lengths a_0 and a_1 related to the polynomial parameters α and β. The figures are in one-to one correspondence to those in Figure 3.*

value, because of the large $O(p^2)$ contribution, tiny dispersive contribution and the fact that within the standard reparametrization of the bare expansion also the second term from the expansion of the denominator is taken into account.

X	Z	r	$10^4 c_{20}$	$10^3 c_{01}$
X^{std}	Z^{std}	r^{std}	-7.10	4.86
X^{std}	Z^{std}	r^{*std}	-6.97	4.79
X^{std}	Z^{std}	r_2	-7.04	4.83

Table 4: *The values of the subthreshold parameters c_{20} and c_{01} related to the polynomial parameters γ and ω at the standard reference point.*

Let us now proceed to the "doubly dangerous" observables a_0 and a_1. These are related to the values of the amplitude at the threshold and receive therefore large contribution from the dispersive part of the amplitude. While a_0 is reproduced well at the standard reference point, a_1 (which starts at the NLO) is off the standard value roughly by a factor 0.6 from the same reasons as for the c_{ij} parameters. The dependence of this observables on X, Z and r is depicted in Fig. 5.

6.4 The role of the remainders

Up to now we have not discussed the uncertainties of the observables calculated within the resummed scheme. They are connected with the direct and indirect remainders as well as with the LEC's L_i, $i = 1, 2, 3$. As the first illustration, we have added the error bars stemming from the remainders to the central values of the various observables depicted in the figures 3-5. These illustrate the rough estimate of the remainders $\delta \sim (30\%)^2 \sim 0.1$ as suggested in [8] and adding the uncertainties in quadrature.

In more detail, at the standard reference point $X = X^{std}$, $Z = Z^{std}$ and $r = r^{std}$, using (5.15, 5.16) and (6.2, 6.13), we numerically get for the corresponding variations (to the first order in the remainders)

$$\begin{aligned} \frac{\Delta\alpha}{\alpha^{CA}} &= (\delta_\alpha + 6.92\delta_{F_\eta M_\eta} - 12.71\delta_{F_K M_K} - 0.76\delta_{F_\pi M_\pi} + 9.37\delta_{F_K} + 5.22\delta_{F_\pi} - 6.92\delta_{F_\eta}) \\ &\quad + (3.38\Delta L_1 + 0.56\Delta L_3) \times 10^3 && (6.18) \\ \Delta\beta &= [(0.69\delta_\beta + 2.34\delta_{F_K} - 20.52\delta_{F_\pi}) + (17.0\Delta L_1 - 2.8\Delta L_3) \times 10^3] \times 10^{-3} M_\eta^2. && (6.19) \end{aligned}$$

This reveals strong sensitivity on both δs and LEC's. Assuming again the typical size of the remainders to be $\delta \sim 0.1$ and adding all the uncertainties in quadrature (for ΔL_i we take the error bars form [44]) we obtain rough (over)estimates

$$\begin{aligned} \left|\overline{\frac{\Delta\alpha}{\alpha^{CA}}}\right| &= \sqrt{1.93^2 + 1.19^2} = 2.27 && (6.20) \\ \left|\overline{\Delta\beta}\right| &= \sqrt{2.06^2 + 5.96^2} \times 10^{-3} M_\eta^2 = 6.31 \times 10^{-3} M_\eta^2, && (6.21) \end{aligned}$$

where the first number under the square root represents the contribution of the remainders while the second accumulates the uncertainty from $L_{1,3}$. Though these numbers are a little bit more optimistic than those in the last row of Table 1 (note that the latter originated purely from the uncertainties of ΔL_i and did not include any estimates of the higher order corrections to α and β), it is clear that, without more restrictive information on the remainders (and L_i, $i = 1, 2, 3$)[25], the predictive power of χPT is reduced considerably in the case of $\pi\eta$ scattering even for "good" observables. In other words, small remainders are not a guarantee of an equivalently small final uncertainty. In what follows, we therefore try to gain some additional information outside the (resummed) χPT expansion to get further estimates of the size of the remainders.

The sources of the remainders are twofold: on one hand there are the unknown terms of the pure derivative expansion, on the other hand the contributions coming from the expansion in the quark masses. We try to get estimates for both of them from different sources, namely using the resonance estimate for the first type as well as independent information from generalized χPT for the second.

6.5 Resonances estimate of the direct remainders

In order to partially estimate the derivative part of the higher order corrections to the chiral expansion, we use the assumption, that the process under consideration is saturated by the exchange of the lowest laying resonances, the interactions of which can be described by the Lagrangian of the resonance chiral theory ($R\chi T$). The leading order Lagrangian of $R\chi T$ was originally formulated in the seminal paper [46] and applied to $\pi\eta$ scattering in [38]. To this process, only scalar resonances as well as $\eta_8 - \eta_0$ mixing contribute. Our result for the amplitude agree with [38] (cf. Appendix J), which we can rewrite in terms of the resonance contribution $G^R_{\pi\eta}$ to $G_{\pi\eta}$ in the form

---

[25] As already discussed, the explicit dependence on these constants could be eliminated by means of reparametrization similar to those for L_i, $i = 4, \ldots, 6$ using further experimental input *e.g.* form K_{e4} decay. The price to pay is to introduce additional remainders connected with observables used for such a reparametrization.

$$G^R_{\pi\eta}(s,t;u) = \alpha_R^{(4)} + \beta_R^{(4)} t + \gamma_R^{(4)} t^2 + \omega_R^{(4)}(s-u)^2 + \Delta G^R_{\pi\eta}(s,t;u). \tag{6.22}$$

The polynomial part with the coefficients (in what follows, M_S and M_{S_1} are the octet and singlet scalar mass respectively, c_d, c_m, $\widetilde{c}_d$, $\widetilde{c}_m$and $\widetilde{d}_m$ are the couplings defined in [46])

$$\begin{aligned}
\alpha_R^{(4)} &= 8M_\pi^2 M_\eta^2 \left(-\frac{c_d^2}{3M_S^2} + \frac{2\widetilde{c}_d^2}{M_{S_1}^2} \right) + 16 M_\pi^2 \overset{o}{M}{}_\eta^{2} \left(\frac{c_d c_m}{3M_S^2} - \frac{\widetilde{c}_d \widetilde{c}_m}{M_{S_1}^2} \right) \\
&\quad + 8 M_\eta^2 \overset{o}{M}{}_\pi^{2} \left(\frac{c_d c_m}{3M_S^2} - \frac{2\widetilde{c}_d \widetilde{c}_m}{M_{S_1}^2} \right) - 16 \frac{\widetilde{d}_m^2}{M_{\eta_1}^2} \overset{o}{M}{}_\pi^{2} \left(\overset{o}{M}{}_\pi^{2} - \overset{o}{M}{}_\eta^{2} \right) \\
&\quad + 20 \overset{o}{M}{}_\pi^{2} \overset{o}{M}{}_\eta^{2} \left(\frac{\widetilde{c}_m^2}{M_{S_1}^2} - \frac{c_m^2}{3M_S^2} \right) - M_\pi^2 \overset{o}{M}{}_\pi^{2} \frac{8 c_d c_m}{3M_S^2} + 4 \overset{o}{M}{}_\pi^{4} \left(\frac{7 c_m^2}{M_S^2} + \frac{\widetilde{c}_m^2}{M_{S_1}^2} \right) &\quad (6.23) \\
\beta_R^{(4)} &= -\frac{8\widetilde{c}_d \widetilde{c}_d}{M_{S_1}^2} \Sigma_{\eta\pi} + \frac{4}{3} \frac{c_d^2}{M_S^2} \Sigma_{\eta\pi} + 8 \left(\frac{\widetilde{c}_d \widetilde{c}_m}{M_{S_1}^2} - \frac{c_d c_m}{3M_S^2} \right) \left(\overset{o}{M}{}_\pi^{2} + \overset{o}{M}{}_\eta^{2} \right) &\quad (6.24) \\
\gamma_R^{(4)} &= \frac{4\widetilde{c}_d^2}{M_{S_1}^2} - \frac{c_d^2}{3M_S^2} &\quad (6.25) \\
\omega_R^{(4)} &= \frac{c_d^2}{3M_S^2} &\quad (6.26)
\end{aligned}$$

gathers the complete $O(p^4)$ resonance contribution (here we can recognize the resonance saturation of the LEC's in (4.23)-(4.26)). This part of the amplitude is already included in our resummed version of χPT, either explicitly through the LEC's $L_1 \ldots L_3$ or implicitly using the reparametrization in terms of the masses, decay constants and parameters r, X and Z. On the other hand, $\Delta G_{\pi\eta,R}(s,t;u)$ can be formally understood as an infinite sum of the higher order corrections in the (purely) derivative expansion, summed up to

$$\begin{aligned}
\Delta G_{\pi\eta,R}(s,t;u) = {} & \frac{4t}{M_{S_1}^2 (M_{S_1}^2 - t)} \left(\widetilde{c}_d (t - 2M_\pi^2) + 2\widetilde{c}_m \overset{o}{M}{}_\pi^{2} \right) \left(\widetilde{c}_d (t - 2M_\eta^2) + 2\widetilde{c}_m \overset{o}{M}{}_\eta^{2} \right) \\
& + \frac{2}{3} \frac{s}{M_S^2 (M_S^2 - s)} \left(c_d (s - M_\pi^2 - M_\eta^2) + 2 c_m \overset{o}{M}{}_\pi^{2} \right)^2 \\
& + \frac{2}{3} \frac{u}{M_S^2 (M_S^2 - u)} \left(c_d (u - M_\pi^2 - M_\eta^2) + 2 c_m \overset{o}{M}{}_\pi^{2} \right)^2 \\
& - \frac{2}{3} \frac{t}{M_S^2 (M_S^2 - t)} \left(c_d (t - 2M_\pi^2) + 2 c_m \overset{o}{M}{}_\pi^{2} \right) \\
& \times \left(c_d (t - 2M_\eta^2) + 2 c_m (2 \overset{o}{M}{}_\eta^{2} - \overset{o}{M}{}_\pi^{2}) \right) \\
& + 16 \frac{\widetilde{d}_m^2 M_\eta^2}{M_{\eta_1}^2 (M_{\eta_1}^2 - M_\eta^2)} \overset{o}{M}{}_\pi^{2} \left(\overset{o}{M}{}_\eta^{2} - \overset{o}{M}{}_\pi^{2} \right).
\end{aligned} \tag{6.27}$$

Of course, this does not exhaust all of the possible higher order corrections (note *e.g.* that the resonance Lagrangian we use contains only the leading order interaction terms with one resonance field and chiral building blocks of the order $O(p^2)$), nevertheless we can use it at least as a rough estimate of the effect of higher orders of the derivative expansion. This is in some sense a procedure opposite to the usual resonance saturation; instead of LEC's we "saturate" the remainders by means of sewing together the resummed chiral expansion $G^{\chi PT}_{\pi\eta}$ (without remainders) with the resonance



Figure 6: *Dependence of the resonance estimates of the direct remainders on $Y = X/Z$ for $r = 15$ (dots), r_2 (solid) and 30 (dashed). Note that for $M_{S_1} = M_S$ and $\widetilde{c}_{m,d} = c_{m,d}/\sqrt{3}$, the remainder δ_β^R is exactly independent on r.*

chiral theory writing the full $R\chi T$ amplitude as

$$G_{\pi\eta}^{R\chi T}(s,t;u) = G_{\pi\eta}^{\chi PT}(s,t;u) + \Delta G_{\pi\eta}^{R}(s,t;u). \tag{6.28}$$

and identifying $G_{\pi\eta}^{R\chi T}$ with the full χPT amplitude, the remainder being $\Delta G_{\pi\eta}^{R}$. Under this assumption, we can derive the following higher order contributions to the direct remainders from $\Delta G_{\pi\eta}^{R}$

$$\Delta\alpha_R = \frac{1}{3}F_\pi^2 M_\pi^2 \delta_\alpha^R = \frac{16}{3}c_m^2 \overset{o}{M}{}_\pi^4 \frac{\Sigma_{\eta\pi}}{M_S^2(M_S^2-\Sigma_{\eta\pi})} + 16\frac{\widetilde{d}_m^2 M_\eta^2}{M_{\eta_1}^2(M_{\eta_1}^2 - M_\eta^2)} \overset{o}{M}{}_\pi^2 \left(\overset{o}{M}{}_\eta^2 - \overset{o}{M}{}_\pi^2\right) \tag{6.29}$$

$$\begin{aligned}\Delta\beta_R = \beta\delta_\beta^R &= \frac{16}{M_{S_1}^4}\left(\widetilde{c}_d M_\pi^2 - \widetilde{c}_m \overset{o}{M}{}_\pi^2\right)\left(\widetilde{c}_d M_\eta^2 - \widetilde{c}_m \overset{o}{M}{}_\eta^2\right)\\ &+\frac{8}{3}\frac{1}{M_S^4}\left(c_d M_\pi^2 - c_m \overset{o}{M}{}_\pi^2\right)\left(c_m(2\overset{o}{M}{}_\eta^2 - \overset{o}{M}{}_\pi^2) - c_d M_\eta^2\right)\\ &-\frac{8}{3}\frac{c_m^2 \overset{o}{M}{}_\pi^4}{M_S^2(M_S^2-\Sigma_{\eta\pi})} - \frac{8}{3}\frac{c_d c_m \overset{o}{M}{}_\pi^2 \Sigma_{\eta\pi}}{M_S^2(M_S^2-\Sigma_{\eta\pi})} - \frac{8}{3}\frac{c_m^2 \overset{o}{M}{}_\pi^4 \Sigma_{\eta\pi}}{M_S^2(M_S^2-\Sigma_{\eta\pi})^2}\end{aligned} \tag{6.30}$$

and similarly for δ_γ^R and δ_ω^R (see Appendix J). Note that the dependence on X and Z is exclusively through the ratio $Y = X/Z$ here.

One can notice that there are two distinct features of this procedure as compared to the usual LEC saturation. First, there is no need to fix a saturation scale, which is the result of "saturating" the renormalization scale independent remainder instead of the scale dependent LEC's. And second, as the resonance contribution are resummed to all chiral orders, the resonance poles are explicitly present in our result, as can be seen in (6.27), (6.29) and (6.30) as well as the formulae for δ_γ^R and δ_ω^R in the Appendix J.

For the rough numerical estimates we use M_S, M_{S_1}, M_{η_1} and the couplings c_d, c_m, $\widetilde{c}_d$, $\widetilde{c}_m$, $\widetilde{d}_m$ from [46]. This gives at the standard reference point $X = X^{std}$, $Z = Z^{std}$ and $r = r^{std}$

$$\delta_\alpha^R = 1.00 \tag{6.31}$$

$$\beta\delta_\beta^R = -0.15 \times 10^{-3} M_\eta^2, \tag{6.32}$$

which represents roughly 55% and 20% correction to values in the first row of the Table 3 respectively. The dependence of δ_α^R and δ_β^R on $Y = X/Z$ and r is depicted in Figure 6. The effect of the resonance remainder estimate on the parameters α and β in a wider range of the X, r and Z is illustrated in the first column of Figure 8, analogous plots for γ and ω are in Figure 7. As can be seen, these results suggests the conclusion that the derivative part of the expansion could in some cases produce higher order remainders with much bigger value than 10%.



Figure 7: *Polynomial parameters γ and ω depending on Y. Horizontal dashed line: standard $O(p^4)$ and central $R\chi PT$ value. The result with resonance remainder estimates is shown by the solid line.*

6.6 Generalized χPT

In the previous subsection we have tried to estimate the contributions to the remainders generated by the derivative expansion. The resulting expressions (6.29) and (6.30) could, however, gather only terms of at most the second order[26] in the quark mass expansion due to the lowest order resonance Lagrangian used. Also, the indirect remainders have not been included in this way as there is no contribution to them in this simplest approach. For the appraisal of the importance of the missing terms we therefore need additional information. One possibility might be to use a resonance Lagrangian with additional terms of higher chiral order suited for saturation of the $O(p^6)$ LEC's [47] and/or to go to the next-to-next-to leading order in the chiral expansion, this is, however, beyond the scope of our paper.

Instead we try to get some flavour of the size of the effect by means of comparison of our previous results with generalized χPT, which was originally designed to handle the badly convergent quark mass expansion in the case $X \ll 1$ and therefore also includes terms which correspond to higher orders in the standard chiral power counting.

In Subsection 5.4, we have already rewritten the generalized expansion of the parameters α and β (as well as that of the masses and decay constants in the Appendices B, I) in the "resummed" form (5.13–5.16) by means of splitting the "standard" remainders into the "nonstandard" extra terms $\delta^{loops}(\mu)$ and $\delta^{CT}(\mu)$ originating in $G\chi PT$ and the unknown part $\delta^{G\chi PT}$. Therefore, neglecting the latter, the sum $\delta^{loops}(\mu) + \delta^{CT}(\mu)$ could be in a sense interpreted as the rough estimate of the contribution to the standard remainders stemming from the higher orders of the quark mass expansion.

While $\delta^{loops}(\mu)$ are known, $\delta^{CT}(\mu)$ depend on the unknown LEC's of the $G\chi PT$ Lagrangian (cf. Appendix F). We therefore set $\delta^{CT}(\mu) = 0$ at a fixed scale μ and by varying this scale in $\delta^{loops}(\mu)$ from $\mu = M_\eta$ to $\mu = M_\rho$ we can get some information on the contribution of the unknown LEC's (note that $\delta^{loops}(\mu) + \delta^{CT}(\mu)$ is renormalization scale independent). We apply this procedure both to the direct and indirect remainders.

The usual way of handling the generalized χPT expansion is to neglect the unknown remainders $\delta^{G\chi PT}$. We can repeat the considerations from the previous subsection and partially appreciate

---

[26] Note that the physical masses in (6.29), (6.30) originate in the derivative expansion.



Figure 8: *Polynomial parameters α and β depending on X and Z for traditional and low value of r. The dotted line shows the central value for $Z = 0.9$, the dashed one is the same for $Z = 0.5$. The error bars correspond to the* 10% *estimates of the remainders. Left column: resonance estimate, filled areas highlight the $O(p^6)$ and higher corrections to the amplitude generated by resonances (lighter for $Z = 0.9$, darker for $Z = 0.5$). Right column: results with combined resonance and $G\chi PT$ estimate of remainders. Filled areas show the scale dependence ($\mu \sim M_\eta - M_\rho$), lighter for $Z = 0.9$ and darker for $Z = 0.5$.*

them using the resonance estimate. In order to avoid double counting, we have to further modify the resonance contribution to the remainder (6.27) subtracting terms of the order $O(p^4)$ within the generalized power counting in the same way as it was done in the previous subsection (c.f. (6.22))

$$\begin{aligned}\Delta G^{G\chi PT}_{\pi\eta,R}(s,t;u) = {} & \Delta G_{\pi\eta,R}(s,t;u) - \frac{16t}{M_{S_1}^4}\widetilde{c}_m^2\,\overset{o}{M}{}_\pi^2\overset{o}{M}{}_\eta^2 - \frac{16}{M_S^4}c_m^2\,\overset{o}{M}{}_\pi^2\,(\overset{o}{M}{}_\pi^2\,(M_\eta^2+M_\pi^2) - \overset{o}{M}{}_\eta^2\,t) \\ & - \frac{16}{M_{\eta_1}^4}\widetilde{d}_m^2\,\overset{o}{M}{}_\pi^2\left(\overset{o}{M}{}_\eta^2 - \overset{o}{M}{}_\pi^2\right)M_\eta^2. \end{aligned} \tag{6.33}$$

This combined $G\chi PT$ and resonance estimate of the remainders is illustrated in Figure 8, the right column shows the result in the case of the polynomial parameters α and β. The effect of the unknown GχPT LEC's is estimated by their scale dependence. The lines closer to the central $R\chi$PT results with neglected remainders are the ones at the scale $\mu = M_\rho$, *i.e.* the constants are set to zero at the usually chosen scale. The filled grey areas then show the change when the LEC's are set to the difference when moving from the scale $\mu = M_\rho$ to M_η. Admittedly, this assigns quite arbitrary numbers to the LEC's, so the uncertainty should be viewed as a rough estimate which can go both ways. The result can be interpreted as being quite consistent with the 10% estimate of the remainders, though clearly exceeding it for some range of the free parameters X, Z and r.

As for the parameters γ and ω, because their contribution in the polynomial expansion is quadratic in the Mandelstam variables, the GχPT estimate does not contribute here.

7. Summary and conclusions

In this paper we have studied the properties of various variants of the chiral expansion on the concrete example of the $\pi\eta$ scattering with special attention to the possible reparametrization in terms of the physical observables. We have tried to illustrate several issues in detail, namely:

- the necessity of carefully choosing a class of "good" observables for which the condition of global convergence is believed to be satisfied in the sense that the $O(p^6)$ and higher remainders are small and under control.
- the necessity to carefully define the bare expansion of "good" observables. Here we have concentrated on the requirements dictated by the exact renormalization scale independence as well as the exact perturbative unitarity. As we have shown, both these requirements can be met by means of sewing together the strict chiral expansion in terms of the LEC's with the dispersive representation for the corresponding Green function. Nevertheless, the resulting bare expansion is not yet defined uniquely; one has to fix the way how to treat the chiral logs and also the $O(p^2)$ amplitudes entering the dispersive integrals. Though the difference is formally of the same order as the remainder itself, we have found that, it might be numerically significant in some region of the free parameters.
- the properties of the standard chiral expansion, based on the potentially "dangerous" reparametrization of the bare expansion implicitly assuming $X, Z \sim 1$ and $r \sim r_2$. In this case we have established strong sensitivity of the observables for $\pi\eta$ scattering on the $O(p^4)$ LEC's; this plagues the standard prediction with large uncertainty. In the case of L_4 and L_6 this also means strong sensitivity to the vacuum fluctuations of the $\overline{s}s$ pairs and therefore to the deviation from the standard scenario with X, $Z \sim 1$. The unusually large absolute values of the NLO corrections as well as large variations achieved for most of the observables (including the "good" ones) when moving from the $O(p^4)$ fit of the LEC's L_i [44] to the $O(p^6)$ based fit [45] might be interpreted as a signal of the importance of the NNLO corrections within the standard chiral expansion. This seems to be also supported by the sensitivity of the NLO contributions to their form when expressed in terms of the physical masses and decay constants (*i.e.* how the $O(p^2)$ relations like *e.g.* the Gell-Mann-Okubo formula are used).

- properties of the “safe” reparametrization and resummation of the vacuum fluctuations. We have confirmed that, for the “good“ observables, the resummed and standard values coincide near the standard reference point $[X^{std}, Z^{std}, r^{std}]$. Under our working hypothesis, which assumes the “good“ observables to be accompanied with small and controllable remainders, this can be interpreted as a consistence of the standard $O(p^4)$ chiral expansion of “good“ observables in the sense that the potentially large higher order remainders are in fact small. On the other hand, in most cases of the “dangerous“ observables the standard and resummed values do not meet at $[X^{std}, Z^{std}, r^{std}]$ (typically for those of them that start at $O(p^4)$). Though this might indicate that the standard expansion is convergent less satisfactorily in this case and the higher order remainders might be important here, the difference between the standard and resummed values lies within the estimated uncertainty of the resummed prediction. Away from the standard reference point, however, we have established that the central values diverge substantially from those of the standard approach even for the “good“ observables. This is a signal that, unless $X, Z \sim 1$, the higher order remainders of the standard chiral expansion might be huge in comparison with LO+NLO value. Though this feature does not exclude the possibility that in this case the standard remainders might be saturated by the NNLO corrections, it could be nevertheless interpreted as an indication of the instability of the standard chiral expansion.

- the role of the remainders within the resummed approach. We have found a strong sensitivity of the observables connected to $\pi\eta$ scattering to the higher order remainders. This might reduce the predictive power of this approach, unless additional information on the actual size of the remainders is available. We have tried to make an independent estimate of the remainders using the simplest version of the resonance chiral Lagrangian as well as making a comparison with $G\chi PT$. Both these estimates seem to be in accord with the rough expectation $\delta \sim 10\%$ for the remainders only in some range of the parameters. For some observables and some corners of the parameter space, they can be substantially larger. Of course, the convergence properties of the bare expansion deserves further investigation by means of going to the NNLO, which is, however, beyond the scope of our article.

As a final remark, we would like to comment on the outlook of future usability and practicality of the outlined procedures in a broader picture. One can easily argue, that even if the $O(p^4)$ standard χPT was considered inadequate, proceeding one step further in the chiral expansion, as was reviewed *e.g.* in [23], is the next logical step to do. Large $O(p^8)$ remainders could alter the precision of the results, but hardly their overall validity. On the other hand pushing the resummation an extra order brings difficulties with the exact algebraic reparametrization.

However, at standard χPT $O(p^6)$ order one encounters the ultimate practical limit of effective theories, the growing number of LEC’s. Additional information beyond core χPT has to be gathered, usually resonance saturation is used. In this sense, the resummed χPT framework in the present form makes a barrier between the explicitly manageable part of the chiral expansion and the remainders which need to be estimated at the right point. If *e.g.* exactly the same RχT Lagrangian to $O(p^6)$ was used, the method of incorporating resonances outlined above would in fact retain more information about the complete RχT theory than usual $O(p^6)$ LEC saturation. Also, explicitly calculable contributions like double logs could be included into remainders similarly to one-loop generalized χPT terms in this paper. Thus, the possible advantages of resumming the vacuum fluctuations could be exploited with only moderate additional computational effort.

Acknowledgment: This work was supported in part by the Center for Particle Physics (project no. LC 527) and by the EU Contract No. MRTN-CT-2006-035482, “FLAVIAnet”.

A. Standard chiral expansion of the parameters X, Z and r

Here we summarize the formulae leading to the standard values of X, Z, Y and r used in Subsection 6.3. Using the standard reparametrization rules explained in Subsection 5.1, we get up to the NLO order in terms of the $O(p^4)$ LEC's

$$\begin{aligned}
X^{std} &= 1 - \frac{M_\pi^2}{2F_\pi^2}\,(32(L_6^r(r_2+2) + L_8^r) \\
&\quad +3J^r_{\pi\pi}(0) + (r_2+1)J^r_{KK}(0) + \frac{1}{9}(2r_2+1)J^r_{\eta\eta}(0) + \frac{11r_2+37}{144\pi^2}\Bigg) \\
Z^{std} &= 1 - \frac{M_\pi^2}{2F_\pi^2}\left(16\,(L_4^r(r_2+2) + L_5^r) + (r_2+1)J^r_{KK}(0) + 4J^r_{\pi\pi}(0) + \frac{r_2+5}{16\pi^2}\right) \\
r^{std} &= r_2 - \frac{M_\pi^2(r_2+1)}{2F_\pi^2}\left(8(2L_8^r - L_5)(r_2-1) - \frac{1}{3}(2r_2+1)J^r_{\eta\eta}(0) + J^r_{\pi\pi}(0) - \frac{(r_2-1)}{24\pi^2}\right).
\end{aligned} \tag{A.1}$$

For r^{std} we can also use an alternative expression based on the chiral expansion of r_2^*

$$\begin{aligned}
r^{*std} &= r_2^* - \frac{M_\pi^2(r_2+1)}{2F_\pi^2}\,(16L_8^r(r_2-1) \\
&\quad +\frac{1}{6}(2r_2+1)J^r_{\eta\eta}(0) + \frac{1}{2}(r_2+1)J^r_{KK}(0) - \frac{3}{2}J^r_{\pi\pi}(0) + \frac{5(r_2-1)}{96\pi^2}\Bigg).
\end{aligned} \tag{A.2}$$

Δ_{GMO} has the following standard chiral expansion

$$\begin{aligned}
\Delta^{std}_{GMO} &= \frac{M_\pi^2(r_2-1)}{2F_\pi^2}\,(32(2L_7^r + L_8^r)(r_2-1) \\
&\quad +\frac{1}{3}(2r_2+1)J^r_{\eta\eta}(0) + (r_2+1)J^r_{KK}(0) - 3J^r_{\pi\pi}(0) + \frac{5(r_2-1)}{48\pi^2}\Bigg).
\end{aligned} \tag{A.3}$$

B. Chiral expansion of the η decay constant

For the bare expansion of the "good" observables F_P^2 we rewrite the standard formulae in the form

$$\begin{aligned}
F_\pi^2 &= F_0^2\left[1 + \frac{B_0\widehat{m}}{F_0^2}\left(16L_4^r(\mu)(r+2) + 16L_5^r(\mu) + (r+1)J^r_{KK}(0) + 4J^r_{\pi\pi}(0) + \frac{1}{16\pi^2}(r+5)\right)\right] \\
&\quad +F_\pi^2\delta_{F_\pi},
\end{aligned} \tag{B.1}$$

$$\begin{aligned}
F_K^2 &= F_0^2\Bigg[1 + \frac{B_0\widehat{m}}{F_0^2}\Bigg(16L_4^r(\mu)(r+2) + 8L_5^r(\mu)(r+1) + \frac{3}{2}(r+1)\left(J^r_{KK}(0) + \frac{1}{16\pi^2}\right) \\
&\quad +\frac{3}{2}\left(J^r_{\pi\pi}(0) + \frac{1}{16\pi^2}\right) + \frac{1}{2}(2r+1)\left(J^r_{\eta\eta}(0) + \frac{1}{16\pi^2}\right)\Bigg)\Bigg] \\
&\quad +F_K^2\delta_{F_K}
\end{aligned} \tag{B.2}$$

$$\begin{aligned}
F_\eta^2 &= F_0^2\left[1 + \frac{B_0\widehat{m}}{F_0^2}\left(16L_4^r(\mu)(r+2) + \frac{16}{3}L_5^r(\mu)(2r+1) + 3(r+1)\left(J^r_{KK}(0) + \frac{1}{16\pi^2}\right)\right)\right] \\
&\quad +F_\eta^2\delta_{F_\eta}.
\end{aligned} \tag{B.3}$$

Within the standard χPT, the $O(p^2)$ parameters B_0, F_0 and r are expressed using inverted expansions of the observables F_P^2, M_P^2, as explained in subsection 5.1. This yields the standard formula for F_η^2

$$\begin{aligned}
F_\eta^2 &= F_\pi^2\left[1 + \frac{M_\pi^2}{F_\pi^2}\left(\frac{16}{3}L_5^r(\mu)(r_2-1) + (r_2+1)J^r_{KK}(0) - 2J^r_{\pi\pi}(0) + \frac{1}{16\pi^2}(r_2-1)\right)\right] \\
&\quad +F_\eta^2\delta^{st}_{F_\eta}.
\end{aligned} \tag{B.4}$$

with a potentially large remainder $\delta^{st}_{F_\eta}$. Numerically, with $L^r_5(M_\rho)$ taken from [44] we get

$$F^2_\eta = 1.625 F^2_\pi. \tag{B.5}$$

On the other hand, the "safe" reparametrization in terms of r, X and Z gives

$$\begin{aligned} F^2_\eta \;=\; & F^2_\pi \left[1 + \frac{2}{3}(r-1)\eta(r) - \frac{1}{3}\frac{M^2_\pi}{F^2_\pi}\left(\frac{X}{Z}\right)\left(J^r_{\pi\pi}(0) - 2(r+1)J^r_{KK}(0) + (2r+1)J^r_{\eta\eta}(0)\right)\right] \\ & + \frac{1}{3}\left(3F^2_\eta\delta_{F_\eta} + F^2_\pi\delta_{F_\pi} - 4F^2_K\delta_{F_K}\right), \end{aligned} \tag{B.6}$$

which is valid as an exact algebraic identity[27].

Following the $G\chi PT$ procedure outlined in Section 5.4, after identifying the corresponding LEC's in both approaches

$$\begin{aligned} \frac{B_0\widehat{m}}{F^2_0}L^r_4(\mu) \;&\to\; \frac{1}{8}\widehat{m}\widetilde{\xi} \\ \frac{B_0\widehat{m}}{F^2_0}L^r_5(\mu) \;&\to\; \frac{1}{8}\widehat{m}\xi^r, \end{aligned} \tag{B.7}$$

and defining the remainders using the physical masses inside the chiral logs, we can use the exact formula (B.3) and write the remainder δ_{F_η} within $G\chi PT$ as

$$F^2_\eta\delta_{F_\eta} = F^2_\eta\delta^{loop}_{F_\eta}(\mu) + F^2_0\delta^{(4)\,CT}_{F_\eta}(\mu) + F^2_\eta\delta^{G\chi PT}_{F_\eta}.$$

Here

$$F^2_\eta\delta^{loop}_{F_\eta}(\mu) = 3\widehat{m}^2\,(r+1)\left(A_0\,(r+1) + 2(r+2)Z^S_0\right)\left(J^r_{KK}(0) + \frac{1}{16\pi^2}\right) \tag{B.8}$$

is the extra loop contribution and $\delta^{(4)\,CT}_{F_\eta}(\mu)$ the contribution of the counterterms from the $O(p^4)$ $G\chi PT$ Lagrangian renormalized at scale μ

$$\begin{aligned} \delta^{(4)\,CT}_{F_\eta}(\mu) \;=\; & \frac{2}{3}\widehat{m}^2\left[\frac{1}{2}(2A_1 + A_2 + 4A_3 + 2B_1 - 2B_2)(1+2r^2)\right. \\ & \left. + 3(A_4 + 2B_4)(1 + \frac{1}{2}r^2) - 4\,C^P_1(r-1)^2 + 2D^S\,(r+2)\,(2r+1)\right]. \end{aligned} \tag{B.9}$$

$\delta^{G\chi PT}_{F_\eta}$ is a new remainder, which is exactly independent on the renormalization scale.

Analogously, for the $G\chi PT$ formula for F^2_π we have, besides the substitution (B.7) to (B.1), to insert

$$F^2_\pi\delta_{F_\pi} = F^2_\pi\delta^{loop}_{F_\pi}(\mu) + F^2_0\delta^{(4)\,CT}_{F_\pi}(\mu) + F^2_\pi\delta^{G\chi PT}_{F_\pi}, \tag{B.10}$$

---

[27] This identity can be also rewritten as

$$4F^2_K(1-\delta_{F_K}) - F^2_\pi(1-\delta_{F_\pi}) - 3F^2_\eta(1-\delta_{F_\eta}) = \left(\frac{X}{Z}\right)M^2_\pi\left(J^r_{\pi\pi}(0) - 2(r+1)J^r_{KK}(0) + (2r+1)J^r_{\eta\eta}(0)\right).$$

Within the standard approach, the parameters on the r.h.s. of this identity can be expressed to the order $O(p^4)$ in terms of the physical observables and it is interpreted as a $O(p^4)$ sum rule

$$4F^2_K - F^2_\pi - 3F^2_\eta = M^2_\pi\left(J^r_{\pi\pi}(0) - 2(r_2+1)J^r_{KK}(0) + (2r_2+1)J^r_{\eta\eta}(0)\right).$$

This gives

$$F^2_\eta = 1.697 F^2_\pi.$$

where the loop and counterterm contribution are now

$$
\begin{aligned}
F_\pi^2 \delta_{F_\pi}^{loop}(\mu) &= 8\widehat{m}^2(A_0 + (r+2)Z_0^S)) \left(J^r_{\pi\pi}(0) + \frac{1}{16\pi^2} \right) \\
&\quad + \widehat{m}^2 \, (r+1) \, (A_0 \, (r+1) + 2(r+2) Z_0^S) \left(J^r_{KK}(0) + \frac{1}{16\pi^2} \right) \\
\delta_{F_\pi}^{(4)\ CT}(\mu) &= 2\widehat{m}^2 \left[A_1 + \frac{1}{2} A_2 + 2A_3 + (A_4 \ + 2B_4)(1 + \frac{1}{2} r^2) \right. \\
&\quad \left. + B_1 - B_2 + 2D^S \, (r+2) \right] . \qquad \text{(B.11)}
\end{aligned}
$$

Finally, we have the expression for F_K^2, where the remainder is replaced with

$$
F_K^2 \delta_{F_K} = F_K^2 \delta_{F_K}^{loop}(\mu) + F_0^2 \delta_{F_K}^{(4)\ CT}(\mu) + F_K^2 \delta_{F_K}^{G\chi PT} \qquad \text{(B.12)}
$$

and the loops and counterterms contribute as

$$
\begin{aligned}
F_K^2 \delta_{F_K}^{loop}(\mu) &= 3\widehat{m}^2(A_0 + (r+2)Z_0^S)) \left(J^r_{\pi\pi}(0) + \frac{1}{16\pi^2} \right) \\
&\quad + \frac{3}{2} \widehat{m}^2 \, (r+1) \, (A_0 \, (r+1) + 2(r+2) Z_0^S) \left(J^r_{KK}(0) + \frac{1}{16\pi^2} \right) \\
&\quad + \widehat{m}^2 (A_0 \left(2r^2 + 1 \right) + 2(r-1)^2 Z_0^P + (r+2)(2r+1) Z_0^S) \left(J^r_{\eta\eta}(0) + \frac{1}{16\pi^2} \right) \\
\delta_{F_K}^{(4)\ CT}(\mu) &= \widehat{m}^2 \left[(A_1 + B_1)(r^2+1) + (A_2 - 2B_2) r + 2(A_4 \ + 2B_4)(1 + \frac{1}{2} r^2) \right. \\
&\quad \left. + 2D^S \, (r+2) \, (r+1) \right] . \qquad \text{(B.13)}
\end{aligned}
$$

In order to reparametrize the $G\chi PT$ bare expansion in terms of the masses and decay constants, we can proceed as follows. Because the exact identity (B.6) is valid independently of the version of χPT, we can also use it in the generalized case, provided we rewrite the remainders according to (B.9, B.10) and (B.12). This step eliminates the LEC's ξ and $\widetilde{\xi}$. Collecting the chiral logs together we have

$$
\begin{aligned}
F_\eta^2 &= F_\pi^2 \left[1 + \frac{2}{3}(r-1)\eta(r) - \frac{1}{3F_\pi^2} \left(\widetilde{M}_\pi^2 \left(J^r_{\pi\pi}(0) + \frac{1}{16\pi^2} \right) \right. \right. \\
&\quad \left. \left. - 4\widetilde{M}_K^2 \left(J^r_{KK}(0) + \frac{1}{16\pi^2} \right) + 3\widetilde{M}_\eta^2 \left(J^r_{\eta\eta}(0) + \frac{1}{16\pi^2} \right) \right) \right] + F_\eta^2 \Delta_{F_\eta}^{G\chi PT}(\mu) \qquad \text{(B.14)}
\end{aligned}
$$

where the $O(p^2)$ masses are given by (5.19) and

$$
F_\eta^2 \Delta_{F_\eta}^{G\chi PT}(\mu) = \frac{1}{3} F_0^2 \left(3\delta_{F_\eta}^{(4)\ CT} + \delta_{F_\pi}^{(4)\ CT} - 4\delta_{F_K}^{(4)\ CT} \right) + \frac{1}{3} \left(3F_\eta^2 \delta_{F_\eta}^{G\chi PT} + F_\pi^2 \delta_{F_\pi}^{G\chi PT} - 4F_K^2 \delta_{F_K}^{G\chi PT} \right) . \qquad \text{(B.15)}
$$

The last step consists of replacing the LEC's F_0, A_0, Z_0^S and Z_0^P with the first term of their expansion in terms of the masses and decay constants as described in Subsection 5.4. This corresponds to a further redefinitions of the generalized remainders.

C. Dispersion representation of the $\pi\eta$ amplitude

For the dispersive representation of the amplitude we need the $S-$ and $T-$channel discontinuities at $O(p^4)$. In the following subsections we give a list of the relevant $O(p^2)$ amplitudes $G^{(2) A_i \to ij}$ and $G^{(2) ij \to A_f}$ and $O(p^4)$ discontinuities $\mathrm{disc} G_0^{ij}$ corresponding to the different intermediate states ij.

C.1 $S-$channel discontinuities at $O(p^4)$

- $\pi\eta$ intermediate state

$$
\begin{aligned}
G^{(2)\pi\eta\to\pi\eta} &= \frac{1}{3}F_0^2 \overset{o}{M}{}_\pi^2 \\
\mathrm{disc}G_0^{\pi\eta}(s) &= 2\frac{\lambda^{1/2}(s, M_\pi^2, M_\eta^2)}{s}\left(\frac{1}{32\pi}\frac{1}{3}\overset{o}{M}{}_\pi^2\right)^2 \frac{F_0^4}{F_\pi^2 F_\eta^2}
\end{aligned}
\tag{C.1}
$$

- $\overline{K}K$ intermediate state

$$
\begin{aligned}
G^{(2)\pi\eta\to\overline{K^0}K^0(K^+K^-)} &= -\frac{\sqrt{3}}{4}F_0^2(s - \frac{1}{3}M_\eta^2 - \frac{1}{3}M_\pi^2 - \frac{2}{3}M_K^2) \\
&\quad + \frac{1}{4\sqrt{3}}F_0^2(2\overset{o}{M}{}_K^2 - \overset{o}{M}{}_\pi^2 - \overset{o}{M}{}_\eta^2) \\
G^{(2)\overline{K^0}K^0(K^+K^-)\to\pi\eta} &= -\frac{\sqrt{3}}{4}F_0^2(s - \frac{1}{3}M_\eta^2 - \frac{1}{3}M_\pi^2 - \frac{2}{3}M_K^2) \\
&\quad + \frac{1}{4\sqrt{3}}F_0^2(2\overset{o}{M}{}_K^2 - \overset{o}{M}{}_\pi^2 - \overset{o}{M}{}_\eta^2) \\
\mathrm{disc}G_0^{\overline{K^0}K^0(K^+K^-)}(s) &= 2\sqrt{1 - \frac{4M_K^2}{s}}\left(\frac{1}{32\pi}\right)^2 \frac{3}{16}[(s - \frac{1}{3}M_\eta^2 - \frac{1}{3}M_\pi^2 - \frac{2}{3}M_K^2) \\
&\quad - \frac{1}{3}(2\overset{o}{M}{}_K^2 - \overset{o}{M}{}_\pi^2 - \overset{o}{M}{}_\eta^2)]^2 \frac{F_0^4}{F_K^4}
\end{aligned}
\tag{C.2}
$$

C.2 $T-$channel discontinuities at $O(p^4)$

- $\pi\pi$ intermediate state

$$
\begin{aligned}
G^{(2)\pi\pi\to\pi\pi,\, I=0} &= F_0^2[(s - \frac{4}{3}M_\pi^2) + \frac{5}{6}\overset{o}{M}{}_\pi^2] \\
\mathrm{disc}G_0^{\pi\pi,\, I=0}(s) &= 2\sigma(s)\left(\frac{1}{32\pi}\right)^2 \frac{1}{3}\overset{o}{M}{}_\pi^2 [(s - \frac{4}{3}M_\pi^2) + \frac{5}{6}\overset{o}{M}{}_\pi^2]\frac{F_0^4}{F_\pi^4}
\end{aligned}
\tag{C.3}
$$

- $\eta\eta$ intermediate state

$$
\begin{aligned}
G^{(2)\eta\eta\to\eta\eta} &= -\frac{1}{3}F_0^2\left(\overset{o}{M}{}_\pi^2 - 4\overset{o}{M}{}_\eta^2\right) \\
\mathrm{disc}G_0^{\eta\eta}(s) &= -2\frac{1}{2}\sqrt{1 - \frac{4M_\eta^2}{s}}\left(\frac{1}{32\pi}\right)^2 \frac{1}{9}\overset{o}{M}{}_\pi^2 \left(\overset{o}{M}{}_\pi^2 - 4\overset{o}{M}{}_\eta^2\right)\frac{F_0^4}{F_\eta^4}
\end{aligned}
\tag{C.4}
$$

- $\overline{K}K$ intermediate state[28]

$$
\begin{aligned}
G^{(2)\pi\pi\to\overline{K^0}K^0(K^+K^-),\, I=0} &= \mp\frac{\sqrt{3}}{4}F_0^2[(s - \frac{2}{3}M_\pi^2 - \frac{2}{3}M_K^2) \\
&\quad + \frac{2}{3}(\overset{o}{M}{}_K^2 + \overset{o}{M}{}_\pi^2)] \\
G^{(2)\overline{K^0}K^0(K^+K^-)\to\eta\eta,\, I=0} &= \pm\frac{1}{4}F_0^2[(3s - 2M_K^2 - 2M_\eta^2)
\end{aligned}
$$

---

[28] Let us note

$$
G^{I=0}(s,t;u) = -\frac{1}{\sqrt{3}}\delta^{ab}G^{ab}(s,t;u) = -\sqrt{3}G(s,t;u)
$$

$$
\begin{aligned}
&+(2\overset{o}{M}{}_\eta^{2}-\frac{2}{3}\overset{o}{M}{}_K^{2})] \\
\mathrm{disc}G_0^{\overline{K^0}K^0(K^+K^-),\,I=0}(s) \;=\;& \sqrt{1-\frac{4M_K^2}{s}}\left(\frac{1}{32\pi}\right)^2\frac{1}{16}[(s-\frac{2}{3}M_\pi^2-\frac{2}{3}M_K^2) \\
&+\frac{2}{3}(\overset{o}{M}{}_K^{2}+\overset{o}{M}{}_\pi^{2})] \\
&\times[(3s-2M_K^2-2M_\eta^2)+(2\overset{o}{M}{}_\eta^{2}-\frac{2}{3}\overset{o}{M}{}_K^{2})]]\frac{F_0^4}{F_K^4} \qquad \text{(C.5)}
\end{aligned}
$$

D. The scalar bubble

In this appendix we summarize the formulae for the scalar bubble, defined as

$$
\begin{aligned}
J_{PQ}(q^2) \;&=\; -\mathrm{i}\int\frac{d^dk}{(2\pi)^d}\frac{1}{(k^2-M_P^2+\mathrm{i}0)((k-q)^2-M_Q^2+\mathrm{i}0)} \\
&=\; -2\lambda_\infty+J_{PQ}^r(q^2). \qquad \text{(D.1)}
\end{aligned}
$$

Here, as usual

$$
\lambda_\infty \;=\; \frac{\mu^{d-4}}{16\pi^2}\left(\frac{1}{d-4}-\frac{1}{2}(\ln 4\pi+\Gamma^{'}(1)+1)\right) \qquad \text{(D.2)}
$$

and $J_{PQ}^r(s)=J_{PQ}^r(0)+\overline{J}_{PQ}(s)$, where

$$
J_{PQ}^r(0)=-\frac{1}{16\pi^2}\frac{M_P^2\ln(M_P^2/\mu^2)-M_Q^2\ln(M_Q^2/\mu^2)}{M_P^2-M_Q^2} \qquad \text{(D.3)}
$$

and $\overline{J}_{PQ}(s)$, sometimes called Chew-Mandelstam function, can be expressed by means of once subtracted dispersion relation as

$$
\overline{J}_{PQ}(s)=\frac{s}{16\pi^2}\int_{(M_P+M_Q)^2}^{\infty}\frac{\mathrm{d}x}{x}\frac{\lambda^{1/2}(x,M_P^2,M_Q^2)}{x}\frac{1}{x-s}. \qquad \text{(D.4)}
$$

The explicit form of $\overline{J}_{PQ}(s)$ reads

$$
\overline{J}_{PQ}(s)=\frac{1}{32\pi^2}\left(2+\frac{\Delta_{PQ}}{s}\ln\frac{M_Q^2}{M_P^2}-\frac{\Sigma_{PQ}}{\Delta_{PQ}}\ln\frac{M_Q^2}{M_P^2}+2\frac{(s-(M_P-M_Q)^2)}{s}\sigma_{PQ}(s)\ln\frac{\sigma_{PQ}(s)-1}{\sigma_{PQ}(s)+1}\right), \qquad \text{(D.5)}
$$

where

$$
\begin{aligned}
\Delta_{PQ} \;&=\; M_P^2-M_Q^2 \\
\Sigma_{PQ} \;&=\; M_P^2+M_Q^2 \\
\sigma_{PQ}(t) \;&=\; \sqrt{\frac{s-(M_P+M_Q)^2}{s-(M_P-M_Q)^2}}=\sqrt{1-\frac{4M_PM_Q}{s-(M_P-M_Q)^2}}\,. \qquad \text{(D.6)}
\end{aligned}
$$

In the limit $M_P\to M_Q$ we get

$$
\begin{aligned}
J_{PP}^r(0) \;&=\; -\frac{1}{16\pi^2}\left(\ln\frac{M_P^2}{\mu^2}+1\right) \\
\overline{J}_{PP}(s) \;&=\; \frac{1}{16\pi^2}\left(2+\sigma_{PP}(s)\ln\frac{\sigma_{PP}(s)-1}{\sigma_{PP}(s)+1}\right). \qquad \text{(D.7)}
\end{aligned}
$$

E. $O(p^4)$ constants L_4-L_8 in terms of masses and decay constants

In this Appendix we summarize the formulae used in the text for the reparametrization of bare expansions of "good" observables. We use the abbreviated notation (5.9–5.12). From the bare expansion of "good" variables F_π^2 and F_K^2 we obtain

$$
\begin{aligned}
4\,\overset{o}{M}{}^2_\pi\, L_4^r(\mu) &= \frac{1}{2}(1-Z-\eta(r))\frac{F_\pi^2}{r+2} \\
&\quad -\frac{M_\pi^2}{4(r+2)(r-1)}\frac{X}{Z}[(4r+1)J^r_{\pi\pi}(0)+(r-2)(r+1)J^r_{KK}(0) \\
&\quad -(2r+1)J^r_{\eta\eta}(0)+\frac{(r+2)(r-1)}{16\pi^2}] \\
&\quad +\frac{2F_K^2\delta_{F_K}-(r+1)F_\pi^2\delta_{F_\pi}}{2(r+2)(r-1)}, \qquad \text{(E.1)}\\
4\,\overset{o}{M}{}^2_\pi\, L_5^r(\mu) &= \frac{1}{2}F_\pi^2\eta(r) \\
&\quad +\frac{M_\pi^2}{4(r-1)}\frac{X}{Z}[5J^r_{\pi\pi}(0)-(r+1)J^r_{KK}(0)-(2r+1)J^r_{\eta\eta}(0)-\frac{3(r-1)}{16\pi^2}] \\
&\quad -\frac{F_K^2\delta_{F_K}-F_\pi^2\delta_{F_\pi}}{(r-1)}. \qquad \text{(E.2)}
\end{aligned}
$$

In the same way, from the expansion of $F_P^2 M_P^2$ we get

$$
\begin{aligned}
4\,\overset{o}{M}{}^4_\pi\, L_6^r(\mu) &= \frac{1}{4}\frac{F_\pi^2 M_\pi^2}{r+2}(1-X-\varepsilon(r)) \\
&\quad -\frac{M_\pi^4}{72(r-1)(r+2)}\left(\frac{X}{Z}\right)^2[27rJ^r_{\pi\pi}(0)+9(r+1)(r-2)J^r_{KK}(0) \\
&\quad +(2r+1)(r-4)J^r_{\eta\eta}(0)+\frac{11(r-1)(r+2)}{16\pi^2}] \qquad \text{(E.3)}\\
&\quad -\frac{F_\pi^2M_\pi^2\delta_{F_\pi M_\pi}[(r+1)^2]-4F_K^2M_K^2\delta_{F_KM_K}}{4(r^2-1)(r+2)} \\
4\,\overset{o}{M}{}^4_\pi\, L_7^r(\mu) &= -\frac{1}{8}F_\pi^2M_\pi^2\left(\varepsilon(r)-\frac{\Delta_{GMO}}{(r-1)^2}\right) \\
&\quad -\frac{3(1+r)F_\eta^2M_\eta^2\delta_{F_\eta M_\eta}+(2r^2+r-1)F_\pi^2M_\pi^2\delta_{F_\pi M_\pi}-8rF_K^2M_K^2\delta_{F_KM_K}}{8(r-1)^2(r+1)} \qquad \text{(E.4)}\\
4\,\overset{o}{M}{}^4_\pi\, L_8^r(\mu) &= \frac{1}{4}F_\pi^2M_\pi^2\varepsilon(r) \\
&\quad +\frac{M_\pi^4}{24(r-1)}\left(\frac{X}{Z}\right)^2[9J^r_{\pi\pi}(0)-3(r+1)J^r_{KK}(0)-(2r+1)J^r_{\eta\eta}(0)-\frac{5(r-1)}{16\pi^2}] \\
&\quad -\frac{2F_K^2M_K^2\delta_{F_KM_K}-(r+1)F_\pi^2M_\pi^2\delta_{F_\pi M_\pi}}{2(r^2-1)}. \qquad \text{(E.5)}
\end{aligned}
$$

F. Lagrangian of $G\chi PT$ to $O(p^4)$

Here we give the traditional form of the $G\chi PT$ Lagrangian. In the following formulae

$$
\begin{aligned}
\chi &= \mathcal{M}+s+\mathrm{i}p \\
\nabla U &= \partial U-\mathrm{i}(v+a)U+\mathrm{i}U(v-a) \\
\chi &= \partial\chi-\mathrm{i}(v+a)\chi+\mathrm{i}\chi(v-a). \qquad \text{(F.1)}
\end{aligned}
$$

Up to the order $O(p^4)$, the Lagrangian can be split into the $O(p^2)$, $O(p^3)$ and $O(p^4)$ parts

$$\mathcal{L} = \mathcal{L}_2 + \mathcal{L}_3 + \mathcal{L}_4, \tag{F.2}$$

where

$$\mathcal{L}_n = \sum_{i+j+k=n} \mathcal{L}^{(i,j,k)} \tag{F.3}$$

and (i,j,k) indicates the number of derivatives, χ sources and powers of B_0 respectively. Then for $O(p^2)$ we get

$$\begin{aligned}
\mathcal{L}^{(2,0,0)} &= \frac{F_0^2}{4}\langle \nabla_\mu U^+ \nabla^\mu U\rangle \\
\mathcal{L}^{(0,1,1)} &= \frac{F_0^2}{2}B_0\langle U^+\chi + \chi^+ U\rangle \\
\mathcal{L}^{(0,2,0)} &= \frac{F_0^2}{4}\left(\overline{A}_0\langle (U^+\chi)^2 + (\chi^+U)^2\rangle\right. \\
&\quad \left. + \overline{Z}_0^S\langle U^+\chi + \chi^+U\rangle^2 + \overline{Z}_0^P\langle U^+\chi - \chi^+U\rangle^2\right).
\end{aligned} \tag{F.4}$$

At the order $O(p^3)$ one has

$$\begin{aligned}
\mathcal{L}^{(2,1,0)} &= \frac{F_0^2}{4}(\overline{\xi}\langle \nabla_\mu U^+\nabla^\mu U(\chi^+U + U^+\chi)\rangle + \overline{\overline{\xi}}\langle \nabla_\mu U^+\nabla^\mu U\rangle\langle \chi^+U + U^+\chi\rangle) \\
\mathcal{L}^{(0,3,0)} &= \frac{F_0^2}{4}(\overline{\rho}_1\langle (\chi^+U)^3 + (U^+\chi)^3\rangle + \overline{\rho}_2\langle (\chi^+U + U^+\chi)\chi^+\chi\rangle \\
&\quad +\overline{\rho}_3\langle (\chi^+U)^2 - (U^+\chi)^2\rangle\langle \chi^+U - U^+\chi\rangle \\
&\quad +\overline{\rho}_4\langle (\chi^+U)^2 + (U^+\chi)^2\rangle\langle \chi^+U + U^+\chi \\
&\quad +\overline{\rho}_5\langle \chi^+U + U^+\chi\rangle\langle \chi^+\chi\rangle + \overline{\rho}_6\langle \chi^+U - U^+\chi\rangle^2\langle \chi^+U + U^+\chi\rangle \\
&\quad +\overline{\rho}_7\langle \chi^+U + U^+\chi\rangle^3) \\
\mathcal{L}^{(2,0,1)} &= \frac{F_0^2B_0}{4}\delta_d^{(1)}\langle \nabla_\mu U^+\nabla^\mu U\rangle \\
\mathcal{L}^{(0,2,1)} &= \frac{F_0^2B_0}{4}\left(\delta^{(1)}\overline{A}_0\langle (U^+\chi)^2 + (\chi^+U)^2\right. \\
&\quad \left. + \delta^{(1)}\overline{Z}_0^S\langle U^+\chi + \chi^+U\rangle^2 + \delta^{(1)}\overline{Z}_0^P\langle U^+\chi - \chi^+U\rangle^2\right) \\
\mathcal{L}^{(0,1,2)} &= \frac{F_0^2}{2}B_0^2\delta_\chi^{(1)}\langle U^+\chi + \chi^+U\rangle.
\end{aligned} \tag{F.5}$$

For the $O(p^4)$ Lagrangian, the building blocks are

$$\begin{aligned}
\mathcal{L}^{(4,0,0)} &= L_1\langle \nabla_\mu U^+\nabla^\mu U\rangle^2 + L_2\langle \nabla_\mu U^+\nabla_\nu U\rangle\langle \nabla^\mu U^+\nabla^\nu U\rangle \\
&\quad +L_3\langle \nabla_\mu U^+\nabla^\mu U\nabla_\nu U^+\nabla^\nu U\rangle \\
&\quad -iL_9\left\langle F_{\mu\nu}^R\nabla^\mu U\nabla^\nu U^+ + F_{\mu\nu}^L\nabla^\mu U^+\nabla^\nu U\right\rangle \\
&\quad +L_{10}\left\langle U^+F_{\mu\nu}^RUF_{\mu\nu}^L\right\rangle + H_1\left\langle F_{\mu\nu}^RF^{R\mu\nu}F_{\alpha\beta}^LF^{L\alpha\beta}\right\rangle \\
\mathcal{L}^{(2,1,1)} &= \frac{F_0^2B_0}{4}(\delta^{(1)}\overline{\xi}\langle \nabla_\mu U^+\nabla^\mu U(\chi^+U + U^+\chi)\rangle + \delta^{(1)}\overline{\overline{\xi}}\langle \nabla_\mu U^+\nabla^\mu U\rangle\langle \chi^+U + U^+\chi\rangle) \\
\mathcal{L}^{(2,0,2)} &= \frac{F_0^2B_0^2}{4}\delta_d^{(2)}\langle \nabla_\mu U^+\nabla^\mu U\rangle \\
\mathcal{L}^{(2,2,0)} &= \frac{F_0^2}{4}\{A_1\langle \nabla_\mu U^+\nabla^\mu U(\chi^+\chi + U^+\chi\chi^+U)\rangle
\end{aligned}$$

$$
\begin{aligned}
&+A_2\langle(\nabla_\mu U^+)U\chi^+(\nabla^\mu U)U^+\chi\rangle\\
&+A_3\,\Big\langle\nabla_\mu U^+U(\chi^+\nabla^\mu\chi-\nabla^\mu\chi^+\chi)+\nabla_\mu UU^+(\chi\nabla^\mu\chi^+-\nabla^\mu\chi\chi^+)\Big\rangle\\
&+A_4\langle\nabla_\mu U^+\nabla^\mu U\rangle\langle\chi^+\chi\rangle\\
&+B_1\langle\nabla_\mu U^+\nabla^\mu U(\chi^+U\chi^+U+U^+\chi U^+\chi)\rangle\\
&+B_2\langle\nabla_\mu U^+\chi\nabla^\mu U^+\chi+\chi^+\nabla^\mu U\chi^+\nabla_\mu U\rangle\\
&+B_4\langle\nabla_\mu U^+\nabla^\mu U\rangle\langle\chi^+U\chi^+U+U^+\chi U^+\chi\rangle\\
&+C_1^S\langle\nabla_\mu U\chi^++\chi\nabla_\mu U^+\rangle\langle\nabla^\mu U\chi^++\chi\nabla^\mu U^+\rangle\\
&+C_2^S\langle\nabla_\mu\chi^+U+U^+\nabla_\mu\chi\rangle\langle\nabla^\mu\chi^+U+U^+\nabla^\mu\chi\rangle\\
&+C_3^S\langle\nabla_\mu\chi^+U+U^+\nabla_\mu\chi\rangle\langle\nabla^\mu U^+\chi+\chi^+\nabla^\mu U\rangle\\
&+C_1^P\langle\nabla_\mu U\chi^+-\chi\nabla_\mu U^+\rangle\langle\nabla^\mu U\chi^+-\chi\nabla^\mu U^+\rangle\\
&+C_2^P\langle\nabla_\mu\chi^+U-U^+\nabla_\mu\chi\rangle\langle\nabla^\mu\chi^+U-U^+\nabla^\mu\chi\rangle\\
&+C_3^P\langle\nabla_\mu\chi^+U-U^+\nabla_\mu\chi\rangle\langle\nabla^\mu U^+\chi-\chi^+\nabla^\mu U\rangle\\
&+D^S\langle\nabla_\mu U^+\nabla^\mu U(\chi^+U+U^+\chi)\rangle\langle\chi^+U+U^+\chi\rangle\\
&+D^P\langle\nabla_\mu U^+\nabla^\mu U(\chi^+U-U^+\chi)\rangle\langle\chi^+U-U^+\chi\rangle\}\\
&+H_2\,\big\langle\nabla_\mu\chi\nabla^\mu\chi^+\big\rangle\\
\mathcal{L}^{(0,4,0)}\;=\;&\frac{F_0^2}{4}\{E_1\langle(\chi^+U)^4+(U^+\chi)^4\rangle\\
&+E_2\langle\chi^+\chi(\chi^+U\chi^+U+U^+\chi U^+\chi)\rangle\\
&+E_3\langle\chi^+\chi U^+\chi\chi^+U\rangle\\
&+F_1^S\langle\chi^+U\chi^+U+U^+\chi U^+\chi\rangle^2\\
&+F_2^S\langle(\chi^+U)^3+(U^+\chi)^3\rangle\langle\chi^+U+U^+\chi\rangle\\
&+F_3^S\langle\chi^+\chi(\chi^+U+U^+\chi)\rangle\langle\chi^+U+U^+\chi\rangle\\
&+F_4^S\langle(\chi^+U)^2+(U^+\chi)^2\rangle\langle\chi^+\chi\rangle\\
&+F_1^P\langle\chi^+U\chi^+U-U^+\chi U^+\chi\rangle^2\\
&+F_2^P\langle(\chi^+U)^3+(U^+\chi)^3\rangle\langle\chi^+U+U^+\chi\rangle\\
&+F_3^P\langle\chi^+\chi(\chi^+U-U^+\chi)\rangle\langle\chi^+U-U^+\chi\rangle\\
&+F_5^{SS}\langle(\chi^+U)^2+(U^+\chi)^2\rangle\langle\chi^+U+U^+\chi\rangle^2\\
&+F_6^{SS}\langle\chi^+\chi\rangle\langle\chi^+U+U^+\chi\rangle^2\\
&+F_5^{SP}\langle(\chi^+U)^2+(U^+\chi)^2\rangle\langle\chi^+U-U^+\chi\rangle\\
&+F_6^{SP}\langle\chi^+\chi\rangle\langle\chi^+U-U^+\chi\rangle^2\\
&+F_7^{SP}\langle(\chi^+U)^2-(U^+\chi)^2\rangle\\
&\times\langle\chi^+U-U^+\chi\rangle\langle\chi^+U+U^+\chi\rangle\\
&+H_3\,\big\langle\chi\chi^+\chi\chi^+\big\rangle+H_4\,\big\langle\chi\chi^+\big\rangle^2\\
\mathcal{L}^{(0,3,1)}\;=\;&\frac{F_0^2B_0}{4}(\delta^{(1)}\overline{\rho}_1\langle(\chi^+U)^3+(U^+\chi)^3\rangle+\delta^{(1)}\overline{\rho}_2\langle(\chi^+U+U^+\chi)\chi^+\chi\rangle\\
&+\delta^{(1)}\overline{\rho}_3\langle(\chi^+U)^2-(U^+\chi)^2\rangle\langle\chi^+U-U^+\chi\rangle\\
&+\delta^{(1)}\overline{\rho}_4\langle(\chi^+U)^2+(U^+\chi)^2\rangle\langle\chi^+U+U^+\chi\\
&+\delta^{(1)}\overline{\rho}_5\langle\chi^+U+U^+\chi\rangle\langle\chi^+\chi\rangle+\delta^{(1)}\overline{\rho}_6\langle\chi^+U-U^+\chi\rangle^2\langle\chi^+U+U^+\chi\rangle\\
&+\delta^{(1)}\overline{\rho}_7\langle\chi^+U+U^+\chi\rangle^3)\\
\mathcal{L}^{(0,2,2)}\;=\;&\frac{F_0^2B_0^2}{4}\Big(\delta^{(2)}\overline{A}_0\langle(U^+\chi)^2+(\chi^+U)^2\rangle
\end{aligned}
$$

$$\begin{aligned}
& + \delta^{(2)}\overline{Z}_0^S \langle U^+\chi + \chi^+ U\rangle^2 + \delta^{(2)}\overline{Z}_0^P \langle U^+\chi - \chi^+ U\rangle^2 \Big) \\
\mathcal{L}^{(0,1,3)} &= \frac{F_0^2}{2} B_0^3 \delta_\chi^{(2)} \langle U^+\chi + \chi^+ U\rangle . \qquad \text{(F.6)}
\end{aligned}$$

In fact, identifying F_0 with the Goldstone boson decay constant and $B_0 = \Sigma/F_0^2$ where $\Sigma = -\langle \overline{u}u\rangle_0$ (in the chiral limit), we have

$$\delta_d^{(i)} = \delta_\chi^{(i)} = 0. \qquad \text{(F.7)}$$

As usual, we can also resume the powers of B_0 already at the Lagrangian level and write

$$\mathcal{L}_n = \sum_{i+j=n} \mathcal{L}^{(i,j)} \qquad \text{(F.8)}$$

with

$$\mathcal{L}^{(i,j)} = \sum_k \mathcal{L}^{(i,j,k)} \qquad \text{(F.9)}$$

and denote

$$\begin{aligned}
A_0 &= \overline{A}_0 + B_0\delta^{(1)}\overline{A}_0 + B_0^2\delta^{(2)}\overline{A}_0 + \ldots \\
Z_0^{S,P} &= \overline{Z}_0^{S,P} + B_0\delta^{(1)}\overline{Z}_0^{S,P} + B_0^2\delta^{(2)}\overline{Z}_0^{S,P} + \ldots \\
\xi &= \overline{\xi} + B_0\delta^{(1)}\overline{\xi} + \ldots \\
\widetilde{\xi} &= \overline{\widetilde{\xi}} + B_0\delta^{(1)}\overline{\widetilde{\xi}} + \ldots \\
\rho_i &= \overline{\rho}_i + B_0\delta^{(1)}\overline{\rho}_i + \ldots , \qquad \text{(F.10)}
\end{aligned}$$

etc.. These LEC's without the bars are then used in the main text. Note that while the $O(p^2)$ parameters $\overline{A}_0$, $\overline{Z}_0^{S,P}$ and the $O(p^3)$ LEC's $\overline{\widetilde{\xi}}$, $\overline{\xi}$ are renormalization scale independent, the renormalized resumed parameters $Z_0^{S,P,r}$, A_0^r and $\widetilde{\xi}^r$, ξ^r run with μ in the same way as $16(B_0/F_0)^2 L_{6-8}^r$ and $8B_0/F_0^2 L_{4,5}$ within the standard χPT.

G. Coefficients of the dispersive part of $G\chi PT$ amplitude

In these formulae as well as in the following two appendices, the masses $\widetilde{M_P^2}$ are the generalized $O(p^2)$ masses given by (5.19).

$$\begin{aligned}
\alpha_{\pi\eta}\widetilde{M_\pi^2} &= 2[\widehat{m}B_0 + 8\widehat{m}^2 A_0 + 2\widehat{m}^2 Z_0^S(5r+4) - 8\widehat{m}^2 Z_0^P(r-1)] \\
\alpha_{\pi\eta K}\widetilde{M_\pi^2} &= 4\widehat{m}^2(r^2-1)(A_0 + 2Z_0^P) \\
\alpha_{\pi\pi}\widetilde{M_\pi^2} &= 2\widehat{m}B_0 + 16\widehat{m}^2 A_0 + 4\widehat{m}^2 Z_0^S(r+8) \\
\alpha_{\eta\eta}(4\widetilde{M_\eta^2} - \widetilde{M_\pi^2}) &= \frac{2}{3}\widehat{m}B_0(1+8r) + \frac{16}{3}\widehat{m}^2 A_0(1+8r^2) \\
&\quad + \frac{4}{3}\widehat{m}^2 Z_0^S(8+41r+32r^2) + \frac{32}{3}\widehat{m}^2 Z_0^P(r-1)(4r-1) \\
(\alpha_{\pi K}-1)\widetilde{M_K}\widetilde{M_\pi} &= 6(A_0 + 2Z_0^S)\widehat{m}^2(r+1) \\
\alpha_{\eta K}(2\widetilde{M_\eta^2} - \frac{2}{3}\widetilde{M_K^2}) &= \frac{2}{3}[\widehat{m}B_0(1+3r) + \widehat{m}^2 A_0(3+10r+19r^2) \\
&\quad + 2\widehat{m}^2 Z_0^S(6+19r+11r^2) + 16\widehat{m}^2 Z_0^P r(r-1)] \qquad \text{(G.1)}
\end{aligned}$$

H. Parameters $\alpha - \omega$ within the generalized χPT

Here we summarize the formulae in terms of the decomposition of the remainders. For the parameter α we write

$$\delta_\alpha = \delta_\alpha^{loop} + 3\frac{\widehat{m}^2 F_0^2}{F_\pi^2 M_\pi^2}\delta_\alpha^{CT}(\mu) + \delta_\alpha^{G\chi PT}. \qquad \text{(H.1)}$$

For the counterterm contribution we get

$$\begin{aligned}
\delta_\alpha^{CT}(\mu) = {} & \frac{1}{3}\widehat{m}\left[81\ \rho_1 + \ \rho_2 + \left(80 - 64\ r - 16\ r^2\right)\ \rho_3\right. \\
& + \left(100 + 64\ r + 34\ r^2\right)\ \rho_4 + \left(2\ +\ r^2\right)\ \rho_5 \\
& \left. + \left(96 - 96\, r\right)\ \rho_6 + \left(144\ + 288\ r + 108\ r^2\right)\ \rho_7\right] \\
& + \frac{8}{3}\left[-(B_1 - B_2)\Sigma_{\pi\eta} + 2D^P M_\pi^2(r-1) - 2C_1^P M_\eta^2(r-1) - \frac{1}{2}D^S[\Sigma_{\pi\eta}(5r+4)]\right. \\
& \left. - 2B_4[3M_\eta^2 + M_\pi^2(2r^2+1)]\right] \\
& + \frac{1}{3}\widehat{m}^2\left[256E_1 + 16E_2 + F_1^P(256 - 256r^2) + F_4^S(32 + 16r^2)\right. \\
& + F_1^S(256 + 320r^2) + F_5^{SP}(192 - 320r + 160r^2 - 32r^3) \\
& + F_2^P(240 - 216r - 24r^3) + F_6^{SP}(32 - 32r + 16r^2 - 16r^3) \\
& + F_3^P(16 - 8r - 8r^3) + F_3^S(16 + 10r + 10r^3) \\
& + F_6^{SS}(32 + 40r + 16r^2 + 20r^3) + F_7^{SP}(384 - 160r - 256r^2 + 32r^3) \\
& \left. + F_2^S(400 + 234r + 74r^3) + F_5^{SS}(576 + 720r + 480r^2 + 168r^3)\right]
\end{aligned} \tag{H.2}$$

and the loops contribute as

$$\begin{aligned}
\frac{1}{3}F_\pi^2 M_\pi^2 \delta_\alpha^{loop} = {} & \frac{1}{3}\left\{[\widetilde{M}_\pi^{\,2}\left(3B_0\widehat{m} + 64A_0\widehat{m}^2 + 2Z_0^S\widehat{m}^2(15r+32) - 8Z_0^P\widehat{m}^2(3r-8)\right)]\right. \\
& \left. -6B_0^2\widehat{m}^2\right\}\left(J^r_{\pi\pi}(0) + \frac{1}{16\pi^2}\right) \\
& + \frac{2}{3}\left\{[\widetilde{M}_K^{\,2}\left(B_0\widehat{m} + 2A_0\widehat{m}^2(r+8) + 2Z_0^S\widehat{m}^2(15r+8) - 8Z_0^P\widehat{m}^2(3r-2)\right)]\right. \\
& \left. -B_0^2\widehat{m}^2(r+1)\right\}\left(J^r_{KK}(0) + \frac{1}{16\pi^2}\right) \\
& \frac{1}{9}\left\{[\widetilde{M}_\eta^2(B_0\widehat{m} + 32A_0\widehat{m}^2 - 16Z_0^P\widehat{m}^2(5r-2) + 2Z_0^S\widehat{m}^2(41r+16)]\right. \\
& \left. -\frac{2}{3}B_0^2\widehat{m}^2(2r+1)\right\}\left(J^r_{\eta\eta}(0) + \frac{1}{16\pi^2}\right) \\
& + \frac{2}{9}\left\{[\widetilde{M}_\pi^{\,2} + 4\widehat{m}^2(3A_0 - 4(r-1)Z_0^P + 2(2r+1)Z_0^S)]^2 - 4B_0^2\widehat{m}^2\right\}J^r_{\pi\eta}(0) \\
& + \frac{3}{4}\left\{[\frac{2}{3}\widetilde{M}_\pi^2 - \frac{8}{3}(r-1)\widehat{m}^2(A_0 + 2Z_0^P)]^2 - \frac{16}{9}B_0^2\widehat{m}^2\right\}J^r_{KK}(0) \\
& + \frac{1}{3}\left\{[\widetilde{M}_\pi^{\,2} + 4\widehat{m}^2(3A_0 - 4(r-1)Z_0^P + 2(2r+1)Z_0^S)]\right. \\
& \left. \times[-2M_\pi^2 + \frac{3}{2}\widetilde{M}_\pi^2 + 10\widehat{m}^2(A_0 + 2Z_0^S)] - 2B_0\widehat{m}(3B_0\widehat{m} - 2M_\pi^2)\right\}J^r_{\pi\pi}(0) \\
& + \frac{2}{9}\left\{[\widetilde{M}_\pi^{\,2} + 4\widehat{m}^2(3A_0 - 4(r-1)Z_0^P + 2(2r+1)Z_0^S)]\right. \\
& \times[\widetilde{M}_\eta^2 - \frac{1}{4}\widetilde{M}_\pi^2 + \widehat{m}^2((8r^2+1)A_0 + 8r(r-1)Z_0^P + 2(2r+1)^2 Z_0^S)] \\
& \left. -\frac{1}{3}B_0^2\widehat{m}^2(8r+1)\right\}J^r_{\eta\eta}(0) \\
& + \frac{1}{8}\left\{[-2M_\pi^2 + 2\widetilde{M}_\pi^2 + 8(r+1)\widehat{m}^2(A_0 + 2Z_0^S))]\right.
\end{aligned}$$

$$\begin{aligned} &\times[-6M_\eta^2 + 6\widetilde{M}_\eta^2 - \frac{8}{3}\widetilde{M}_K^2 + \frac{8}{3}(r+1)\widehat{m}^2(3rA_0 + 2(r-1)Z_0^P + 2(2r+1)Z_0^S] \\ &-2(2B_0\widehat{m} - M_\pi^2)(\frac{4}{3}B_0\widehat{m}(4r+1) - 6M_\eta^2)\Big\}\, J^r_{KK}(0). \end{aligned} \tag{H.3}$$

In the same way we have for β

$$\beta\delta_\beta = \beta\delta_\beta^{loop} + \widehat{m}^2 F_0^2 \delta_\beta^{CT}(\mu) + \beta\delta_\beta^{G\chi PT}, \tag{H.4}$$

where

$$\delta_\beta^{CT}(\mu) \;=\; \frac{8}{3}[(C_1^S + D^S)(2r+1) + 2B_4(r^2+2)] \tag{H.5}$$

$$\begin{aligned} \beta\delta_\beta^{loop} \;=\; &-\frac{3}{4}\Big\{[\frac{2}{3}\widetilde{M}_\pi^2 - \frac{8}{3}(r-1)\widehat{m}^2(A_0 + 2Z_0^P)] - \frac{4}{3}B_0\widehat{m}\Big\}\, J^r_{KK}(0) \\ &+\frac{1}{3}\Big\{[\widetilde{M}_\pi^{\,2} + 4\widehat{m}^2(3A_0 - 4(r-1)Z_0^P + 2(2r+1)Z_0^S)] - 2B_0\widehat{m}\Big\}\, J^r_{\pi\pi}(0) \\ &+\frac{1}{8}\Big\{[6(\widetilde{M}_\eta^2 - M_\eta^2 + \widetilde{M}_\pi^2 - M_\pi^2) - \frac{8}{3}\widetilde{M}_K^2 \\ &+\frac{8}{3}(r+1)\widehat{m}^2(3A_0(r+3) + 4Z_0^S(r+5) + 2(r-1)Z_0^P)] \\ &-[\frac{8}{3}B_0\widehat{m}(2r+5) - 6M_\eta^2 - 6M_\pi^2]\Big\}\, J^r_{KK}(0). \end{aligned} \tag{H.6}$$

For the remaining two parameters the corresponding decomposition of the remainders

$$\gamma\delta_\beta(\mu) \;=\; \gamma\delta_\gamma{}^{loops}(\mu) + \widehat{m}^2 F_0^2 \delta_\gamma^{CT}(\mu) + \gamma\delta_\gamma^{G\chi PT} \tag{H.7}$$

$$\omega\delta_\omega(\mu) \;=\; \omega\delta_\omega{}^{loops}(\mu) + \widehat{m}^2 F_0^2 \delta_\omega^{CT}(\mu) + \omega\delta_\omega^{G\chi PT} \tag{H.8}$$

is trivial, *i.e.*

$$\delta_\gamma{}^{loops}(\mu) = \delta_\gamma^{CT}(\mu) = \delta_\omega{}^{loops}(\mu) = \delta_\omega^{CT}(\mu) = 0. \tag{H.9}$$

I. Generalized χPT contributions to the bare expansion remainders for the masses

The expressions for ξ, $\widetilde{\xi}$ can be obtained from the exact algebraic identities (E.2) after identification (5.24) and using the representation (B.10) and (B.11) for the remainder of F_π^2 and (B.12) and (B.13) for the remainder of F_K^2. In the same spirit, A_0, Z_0^S and Z_0^P can be expressed using identities (E.5) and the following remainders

$$\begin{aligned} F_\pi^2 M_\pi^2 \delta_{F_\pi M_\pi} \;&=\; F_\pi^2 M_\pi^2 \delta^{loop}_{F_\pi M_\pi}(\mu) + F_0^2\widehat{m}^2\delta^{CT}_{F_\pi M_\pi}(\mu) + F_\pi^2 M_\pi^2 \delta^{G\chi PT}_{F_\pi M_\pi} \\ F_K^2 M_K^2 \delta_{F_K M_K} \;&=\; F_K^2 M_K^2 \delta^{loop}_{F_K M_K}(\mu) + F_0^2\widehat{m}^2\delta^{CT}_{F_K M_K}(\mu) + F_K^2 M_K^2 \delta^{G\chi PT}_{F_K M_K} \\ F_\eta^2 M_\eta^2 \delta_{F_\eta M_\eta} \;&=\; F_\eta^2 M_\eta^2 \delta^{loop}_{F_\eta M_\eta}(\mu) + F_0^2\widehat{m}^2\delta^{CT}_{F_\eta M_\eta}(\mu) + F_\eta^2 M_\eta^2 \delta^{G\chi PT}_{F_\eta M_\eta}, \end{aligned} \tag{I.1}$$

where

$$
\begin{aligned}
F_\pi^2 M_\pi^2 \delta^{loop}_{F_\pi M_\pi}(\mu) \;=\;& \Big[\widetilde{M}_\pi^2 \big(3B_0\widehat{m} + 16A_0\widehat{m}^2 + 2Z_0^S\widehat{m}^2(3r+16)\big) - 6B_0^2\widehat{m}^2\Big] \\
& \times \left(J^r_{\pi\pi}(0) + \frac{1}{16\pi^2}\right) \\
& +2\Big[\widetilde{M}_K^2 \big(B_0\widehat{m} + 2A_0\widehat{m}^2(r+2) + 2Z_0^S\widehat{m}^2(3r+4)\big) - B_0^2\widehat{m}^2(r+1)\Big] \\
& \times \left(J^r_{KK}(0) + \frac{1}{16\pi^2}\right) \\
& +\frac{1}{3}\Big[\widetilde{M}_\eta^2 (B_0\widehat{m} + 8A_0\widehat{m}^2 - 8Z_0^P\widehat{m}^2(r-1) + 2Z_0^S\widehat{m}^2(5r+4) \\
& -\frac{2}{3}B_0^2\widehat{m}^2(2r+1)\Big] \left(J^r_{\eta\eta}(0) + \frac{1}{16\pi^2}\right)
\end{aligned}
\tag{I.2}
$$

$$
\begin{aligned}
\delta^{CT}_{F_\pi M_\pi}(\mu) \;=\;& \widehat{m}[9\rho_1 + \rho_2 + 2\rho_4(10+4r+r^2) \\
& +\rho_5(2+r^2) + 12\rho_7(4+4r+r^2)] \\
& +2\widehat{m}^2[8E_1 + 2E_2 + 8F_1^S\left(2+r^2\right) + F_2^S\left(9r+r^3+20\right) + F_3^S\left(4+r+r^3\right) \\
& +2F_4^S\left(2+r^2\right) + 4F_5^{SS}\left(r+2\right)\left(r^2+2r+6\right) + 2F_6^{SS}\left(r+2\right)\left(2+r^2\right)],
\end{aligned}
\tag{I.3}
$$

then

$$
\begin{aligned}
F_K^2 M_K^2 \delta^{loop}_{F_K M_K}(\mu) \;=\;& \Bigg\{\frac{3}{4}\Big[\widetilde{M}_\pi^2 \big(B_0\widehat{m} + A_0\widehat{m}^2(r+5) + 2Z_0^S\widehat{m}^2(r+6)\big) - 2B_0^2\widehat{m}^2\Big] \\
& \times \left(J^r_{\pi\pi}(0) + \frac{1}{16\pi^2}\right) \\
& +\frac{3}{2}\Big[\widetilde{M}_K^2 \big(B_0\widehat{m} + 3A_0\widehat{m}^2(r+1) + 2Z_0^S\widehat{m}^2(3r+4)\big) - B_0^2\widehat{m}^2(r+1)\Big] \\
& \times \left(J^r_{KK}(0) + \frac{1}{16\pi^2}\right) \\
& +\frac{1}{12}\Big[\widetilde{M}_\eta^2 \big(5B_0\widehat{m} + A_0\widehat{m}^2(17r+5) + 8Z_0^P\widehat{m}^2(r-1) + 2Z_0^S\widehat{m}^2(13r+14)\big) \\
& -\frac{10}{3}B_0^2\widehat{m}^2\left(2r+1\right)\Big] \left(J^r_{\eta\eta}(0) + \frac{1}{16\pi^2}\right)\Bigg\}\left(r+1\right)
\end{aligned}
\tag{I.4}
$$

$$
\begin{aligned}
\delta^{CT}_{F_K M_K}(\mu) \;=\;& \frac{1}{2}\widehat{m}[3\rho_1(1+r)(1+r+r^2) + \rho_2(1+r^3) + 6\rho_4\left(r+1\right)\left(2+2r+r^2\right) \\
& +\rho_5\left(r+1\right)\left(2+r^2\right) + 12\rho_7\left(r+1\right)\left(r+2\right)^2] \\
& +\widehat{m}^2[2E_1\left(1+r\right)^2\left(1+r^2\right) + E_2\left(1+r\right)^2\left(1-r+r^2\right) \\
& +\frac{1}{2}E_3\left(r^2-1\right)^2 \\
& +4F_1^S\left(1+r\right)^2\left(2+r^2\right) + F_2^S\left(1+r\right)\left(8+9r+9r^2+4r^3\right) \\
& -F_3^S\left(1+r\right)\left(4-r+r^2+2r^3\right) + F_4^S\left(1+r\right)^2\left(2+r^2\right) \\
& +4F_5^{SS}\left(1+r\right)\left(2+r\right)\left(4+3r+2r^2\right) \\
& +2F_6^{SS}\left(1+r\right)\left(2+r\right)\left(2+r^2\right)]
\end{aligned}
\tag{I.5}
$$

and

$$
\begin{aligned}
F_\eta^2 M_\eta^2 \delta^{loop}_{F_\eta M_\eta}(\mu) \;=\;& \Big[\widetilde{M}_\pi^2 \big(B_0\widehat{m} + 8\,A_0\,\widehat{m}^2 - 8\,\widehat{m}^2\,Z_0^P\,(r-1)\, + 2\,\widehat{m}^2\,Z_0^S\,(4+5\,r)\,\big) - 2B_0^2\widehat{m}^2\Big] \\
& \times \left(J^r_{\pi\pi}(0) + \frac{1}{16\pi^2}\right)
\end{aligned}
$$

$$
\begin{aligned}
&+\frac{2}{3}\,\Big[\widetilde{M}_K^2\,\left(B_0\widehat{m}\,\left(1+4\,r\right)+2\,A_0\,\widehat{m}^2\,\left(2+r+8\,r^2\right)+8\,\widehat{m}^2\,Z_0^P\,\left(r-1\right)\,\left(2\,r-1\right)\right.\\
&\left.+2\,\widehat{m}^2\,Z_0^S\,\left(4+15\,r+8\,r^2\right)\right)-B_0^2\widehat{m}^2\left(4\,r+1\right)\left(r+1\right)\Big]\\
&\times\left(J_{KK}^r(0)+\frac{1}{16\pi^2}\right)\\
&+\frac{1}{9}\,\Big[\widetilde{M}_\eta^2\left(B_0\widehat{m}\,\left(1+8\,r\right)+8A_0\,\widehat{m}^2\,\left(1+8\,r^2\right)+16\,\widehat{m}^2\,Z_0^P\,\left(r-1\right)\,\left(4\,r-1\right)\right.\\
&\left.+2\,\widehat{m}^2\,Z_0^S\,\left(8+41\,r+32\,r^2\right)\right)-\frac{2}{3}B_0^2\widehat{m}^2\left(8\,r+1\right)\left(2r+1\right)\Big]\\
&\times\left(J_{\eta\eta}^r(0)+\frac{1}{16\pi^2}\right) \qquad\qquad (\mathrm{I.6})
\end{aligned}
$$

$$
\begin{aligned}
\delta_{F_\eta M_\eta}^{CT}(\mu) \;=\;& \frac{1}{3}\widehat{m}[9\rho_1\left(1+2r^3\right)+\rho_2\left(1+2r^3\right)+16\rho_3\left(r-1\right)^2\left(1+r\right)\\
&+2\rho_4\left(10+8r+17r^2+10r^3\right)\\
&+\rho_5\left(1+2r\right)\left(2+r^2\right)+16\rho_6\left(2-3r+r^3\right)+12\rho_7\left(2+r\right)^2\left(1+2r\right)]\\
&+\frac{2}{3}\widehat{m}^2[8E_1\left(1+2r^4\right)+2E_2\left(1+2r^4\right)\\
&+8F_1^S\left(r^2+2\right)\left(2r^2+1\right)+16F_1^P\left(r^2-1\right)^2\\
&+F_2^S\left(20+13r+37r^3+20r^4\right)+12F_2^P\left(r^2+r+1\right)\left(r-1\right)^2\\
&+F_3^S\left(4+5r+5r^3+4r^4\right)+4F_3^P\left(r^2+r+1\right)\left(r-1\right)^2\\
&+2F_4^S\left(r^2+2\right)\left(2r^2+1\right)\\
&+12F_5^{SS}\left(r+2\right)\left(2r^3+3r^2+2r+2\right)+8F_5^{SP}\left(r^2+2\right)\left(r-1\right)^2\\
&+2F_6^{SS}\left(2r+1\right)\left(r+2\right)\left(r^2+2\right)+4F_6^{SP}\left(r^2+2\right)\left(r-1\right)^2\\
&+16F_7^{SP}\left(r+2\right)\left(r+1\right)\left(r-1\right)^2]. \qquad\qquad (\mathrm{I.7})
\end{aligned}
$$

J. Resonance amplitude and remainders estimates

Here we give the contribution to the amplitude [38] related to the resonance exchange, derived from the leading order Lagrangian of $R\chi T$ (we have confirmed this expression by independent calculation)

$$
\begin{aligned}
G_R(s,t;u) \;=\;& 4\frac{1}{M_{S_1}^2-t}\left(\widetilde{c}_d(t-2M_\pi^2)+2\widetilde{c}_m\,\overset{o}{M}{}_\pi^{2}\right)\left(\widetilde{c}_d(t-2M_\eta^2)+2\widetilde{c}_m\,\overset{o}{M}{}_\eta^{2}\right)\\
&+4\frac{\widetilde{c}_m^2}{M_{S_1}^2}\,\overset{o}{M}{}_\pi^{2}\left(\overset{o}{M}{}_\pi^{2}+\overset{o}{M}{}_\eta^{2}\right)+4\frac{c_m^2}{3M_S^2}\,\overset{o}{M}{}_\pi^{2}\left(\overset{o}{M}{}_\pi^{2}-\overset{o}{M}{}_\eta^{2}\right)\\
&+\frac{2}{3}\frac{1}{M_S^2-s}\left(c_d(s-M_\pi^2-M_\eta^2)+2c_m\,\overset{o}{M}{}_\pi^{2}\right)^2\\
&+\frac{2}{3}\frac{1}{M_S^2-u}\left(c_d(u-M_\pi^2-M_\eta^2)+2c_m\,\overset{o}{M}{}_\pi^{2}\right)^2\\
&-\frac{2}{3}\frac{1}{M_S^2-t}\left(c_d(t-2M_\pi^2)+2c_m\,\overset{o}{M}{}_\pi^{2}\right)\left(c_d(t-2M_\eta^2)+2c_m(2\,\overset{o}{M}{}_\eta^{2}-\overset{o}{M}{}_\pi^{2})\right)\\
&-16\frac{\widetilde{d}_m^2}{M_{\eta_1}^2}\,\overset{o}{M}{}_\pi^{2}\left(\overset{o}{M}{}_\pi^{2}-\overset{o}{M}{}_\eta^{2}\right). \qquad\qquad (\mathrm{J.1})
\end{aligned}
$$

The resonance estimate of the remainders δ_γ^R and δ_ω^R are

$$
\begin{aligned}
\gamma\delta_\gamma^R = & -\frac{8}{3M_S^6}\left(c_d M_\pi^2 - c_m \overset{o}{M}{}_\pi^2\right)\left(c_d M_\eta^2 - c_m(2\overset{o}{M}{}_\eta^2 - \overset{o}{M}{}_\pi^2)\right) \\
& + \frac{4}{3}\frac{c_d}{M_S^4}\left(c_d M_\eta^2 - c_m(2\overset{o}{M}{}_\eta^2 - \overset{o}{M}{}_\pi^2)\right) + \frac{1}{3}\frac{c_d^2\Sigma_{\pi\eta}}{M_S^2(M_S^2 - \Sigma_{\pi\eta})} \\
& + \frac{4}{3}\frac{c_d c_m}{(M_S^2 - \Sigma_{\pi\eta})^2}\overset{o}{M}{}_\pi^2 + \frac{4}{3}\frac{c_m^2}{(M_S^2 - \Sigma_{\pi\eta})^3}\overset{o}{M}{}_\pi^4 \\
& + \frac{16}{M_{S_1}^6}\left(\widetilde{c}_d M_\pi^2 - \widetilde{c}_m \overset{o}{M}{}_\pi^2\right)\left(\widetilde{c}_d M_\eta^2 - \widetilde{c}_m \overset{o}{M}{}_\eta^2\right) \\
& - \frac{8\widetilde{c}_d}{M_{S_1}^4}\left(\widetilde{c}_d\Sigma_{\pi\eta} - \widetilde{c}_m(\overset{o}{M}{}_\pi^2 + \overset{o}{M}{}_\eta^2)\right)
\end{aligned}
\tag{J.2}
$$

$$
\omega\delta_\omega^R = -\frac{1}{3}\frac{c_d^2\Sigma_{\pi\eta}}{M_S^2(M_S^2 - \Sigma_{\pi\eta})} + \frac{4}{3}\frac{c_m^2\overset{o}{M}{}_\pi^4}{(M_S^2 - \Sigma_{\pi\eta})^3} + \frac{4}{3}\frac{c_d c_m}{(M_S^2 - \Sigma_{\pi\eta})^2}\overset{o}{M}{}_\pi^2\,. \tag{J.3}
$$

References

[1] S. Weinberg, Physica A **96** (1979) 327.

[2] J. Gasser and H. Leutwyler, Annals Phys. **158** (1984) 142.

[3] J. Gasser and H. Leutwyler, Nucl. Phys. B **250** (1985) 465.

[4] S. Descotes-Genon, L. Girlanda and J. Stern, JHEP **0001** (2000) 041 [arXiv:hep-ph/9910537].

[5] S. Descotes-Genon and J. Stern, Phys. Rev. D **62** (2000) 054011 [arXiv:hep-ph/9912234].

[6] S. Descotes-Genon and J. Stern, Phys. Lett. B **488** (2000) 274 [arXiv:hep-ph/0007082].

[7] S. Descotes-Genon, L. Girlanda and J. Stern, Eur. Phys. J. C **27** (2003) 115 [arXiv:hep-ph/0207337].

[8] S. Descotes-Genon, N. H. Fuchs, L. Girlanda and J. Stern, Eur. Phys. J. C **34** (2004) 201 [arXiv:hep-ph/0311120].

[9] S. Descotes-Genon, arXiv:hep-ph/0703154.

[10] T. Appelquist, J. Terning, L.C.R. Wijewardhana, Phys. Rev. Lett. **77** (1996), [arXiv:hep-ph/9602385]

[11] T. Appelquist, A. Ratnaweera, J. Terning, L. C. R. Wijewardhana, Phys. Rev. D **58** (1998) 105017, [arXiv:hep-ph/9806472]

[12] E. Gardi, G. Grunberg, JHEP **9903** (1999) 024, [arXiv:hep-th/9810192]

[13] M. Velkovsky, E. Shuryak, Phys. Lett. B **437** (1998) 398, [arXiv:hep-ph/9703345]

[14] Ch.S. Fischer, R. Alkofer, Phys. Rev. D **67** (2003) 094020, [arXiv:hep-ph/0301094]

[15] R.D. Mawhinney, Nucl. Phys. Proc. Suppl. **60A** (1998) 306, hep-lat/9705031

[16] R.D. Mawhinney, Nucl. Phys. Proc. Suppl. **83** (2000) 57, [arXiv:hep-lat/0001032]

[17] Chengzhong Sui, Nucl. Phys. Proc. Suppl. **73** (1999) 228, [arXiv:hep-lat/9811011]

[18] Y. Iwasaki, K. Kanaya, S. Kaya, S. Sakai, T. Yoshie, Prog. Theor. Phys. Suppl. **131** (1998) 415, [arXiv:hep-lat/9804005]

[19] Y. Iwasaki, K. Kanaya, S. Kaya, S. Sakai, T. Yoshie, Phys. Rev. D **69** (2004) 014507, [arXiv:hep-lat/0309159]

[20] L. Girlanda, J. Stern and P. Talavera, Phys. Rev. Lett. **86** (2001) 5858 [arXiv:hep-ph/0103221].

[21] S. Pislak *et al.* [BNL-E865 Collaboration], Phys. Rev. Lett. **87** (2001) 221801 [arXiv:hep-ex/0106071].

[22] S. Descotes-Genon, N. H. Fuchs, L. Girlanda and J. Stern, Eur. Phys. J. C **24** (2002) 469 [arXiv:hep-ph/0112088].

[23] J. Bijnens, Prog. Part. Nucl. Phys. **58** (2007) 521-586 [arXiv:hep-ph/0604043].

[24] N. H. Fuchs, H. Sazdjian and J. Stern, Phys. Lett. B **269** (1991) 183.

[25] J. Stern, H. Sazdjian and N. H. Fuchs, Phys. Rev. D **47** (1993) 3814 [arXiv:hep-ph/9301244].

[26] M. Knecht, B. Moussallam and J. Stern, Nucl. Phys. B **429** (1994) 125 [arXiv:hep-ph/9402318].

[27] M. Knecht, B. Moussallam and J. Stern, *Contribution to the second edition of the DAPHNE physics handbook, L. Maiani, G. Pancheri and N. Paver Eds.*, arXiv:hep-ph/9411259.

[28] M. Knecht and J. Stern, *Contribution to the second edition of the DAPHNE physics handbook, L. Maiani, G. Pancheri and N. Paver Eds.*, arXiv:hep-ph/9411253.

[29] M. Knecht, B. Moussallam, J. Stern and N. H. Fuchs, Nucl. Phys. B **457** (1995) 513 [arXiv:hep-ph/9507319].

[30] B. Moussallam, Eur. Phys. J. C **14** (2000) 111 [arXiv:hep-ph/9909292].

[31] B. Moussallam, JHEP **0008** (2000) 005 [arXiv:hep-ph/0005245].

[32] B. Ananthanarayan, P. Buettiker and B. Moussallam, Eur. Phys. J. C **22** (2001) 133 [arXiv:hep-ph/0106230].

[33] S. Descotes-Genon, JHEP **0103** (2001) 002 [arXiv:hep-ph/0012221].

[34] J. Novotny and M. Kolesar, *Presented at International Conference on High-Energy Interactions: Theory and Experiment (Hadron Structure '02), Herlany, Slovakia, 22-27 Sep 2002.* arXiv:hep-ph/0212311.

[35] M. Kolesar and J. Novotny, *Presented at International Conference on High-Energy Interactions: Theory and Experiment (Hadron Structure '02), Herlany, Slovakia, 22-27 Sep 2002.* arXiv:hep-ph/0301005.

[36] M. Kolesar and J. Novotny, Czech. J. Phys. **54** (2004) B63 [arXiv:hep-ph/0402239].

[37] M. Kolesar and J. Novotny, in preparation

[38] V. Bernard, N. Kaiser and U. G. Meissner, Phys. Rev. D **44** (1991) 3698.

[39] B. Ananthanarayan and P. Buettiker, Eur. Phys. J. C **19** (2001) 517 [arXiv:hep-ph/0012023].

[40] M. Zdráhal and J. Novotný, in preparation

[41] A. Gomez Nicola and J. R. Pelaez, Phys. Rev. D **65** (2002) 054009 [arXiv:hep-ph/0109056].

[42] H. Osborn, Nucl. Phys. B **15** (1970) 501.

[43] M. Kolesar and J. Novotny, *Presented at Hadron Structure'07, Modra, Slovakia*, to be published in Fizika B (Zagreb, Croatia), arXiv:0802.1151

[44] J. Bijnens, G. Ecker and J. Gasser, *Contribution to the second edition of the DAPHNE physics handbook, L. Maiani, G. Pancheri and N. Paver Eds.* arXiv:hep-ph/9411232.

[45] G. Amoros, J. Bijnens and P. Talavera, Nucl. Phys. B **585** (2000) 293 [Erratum-ibid. B **598** (2001) 665] [arXiv:hep-ph/0003258].

[46] G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B **321** (1989) 311.

[47] V. Cirigliano, G. Ecker, M. Eidemuller, R. Kaiser, A. Pich and J. Portoles, Nucl. Phys. B **753** (2006) 139 [arXiv:hep-ph/0603205].